


Onychomycosis Treatment & the Antifungal Drug Chart

(Chart Pages 1 & 2 printed; 3rd page available online)

April 2010



Recent Guidelines:

Canadian : 
Bugs and drugs 2006
<http://www.bugsanddrugs.ca/>

American :
IDSA *Candida* guidelines 2009
<http://www.journals.uchicago.edu/doi/pdf/10.1086/596757>

UK Guideline 2003
<http://bad.org.uk/Portals/Bad/Guidelines/Clinical%20Guidelines/Onychomycosis.pdf>¹

Review Articles:

NEJM: Fungal nail disease 2009
<http://content.nejm.org/cgi/reprint/360/20/2108.pdf>²

Cochrane: Topical fungal treatments of the skin & foot 2007
<http://mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD001434/pdf/fs.html>³

Other Resources:

Images of skin diseases, includes other dermatologic links: www.dermnet.com

Patient Resources:

BMJ Clinical Evidence
http://clinicalevidence.bmj.com/ceweb/conditions/skd/1715/fungal-toenail-infections-standard-ce_patient_leaflet.pdf

Highlights: ★

- 1) Not all abnormal nails are fungal, treat only if culture positive for dermatophyte
- 2) To minimize potential for false negative, culture nail clipping and deep scrapings
- 3) Treat with terbinafine for 12-16 weeks (drug of choice for toenail onychomycosis)
- 4) Mark nail at end of treatment to monitor treatment success

RxFiles Related:

Antifungal chart:
<http://www.rxfiles.ca/rxfiles/uploads/documents/members/cht-antifungal.pdf>

Topical Steroid Chart:
<http://www.rxfiles.ca/rxfiles/uploads/documents/members/CHT-SteroidClassPotencyCOLOR.pdf>

OTC Chart: Fungal Infections
<http://www.rxfiles.ca/rxfiles/uploads/documents/members/CHT-OTCs.pdf>

RxFiles Academic Detailing
Saskatoon City Hospital
Saskatoon, SK Canada

see www.RxFiles.ca

General Overview – Onychomycosis^{4,5,6,7,8}

- Onychomycosis is a fungal infection of the nails most commonly caused by dermatophytes. Less often *Candida* and molds may affect the nail.
- Onychomycosis is recognized by thickening of the distal end of the nail associated with some loosening of the nail plate from the nail bed. The nail plate shows *butter yellow* coloured, vertical bands starting at the distal end of the nail.
- Both toenails and finger nails may be affected, but dermatophyte infections of fingers seldom occur in the absence of toenail infections.
- Fungal infections of the foot are not life-threatening but can cause discomfort and become unsightly. For some, they predispose to recurrent cellulitis of the legs.

Case discussion

- Mr. T., a 69 yr old man reports that his big toenail has some yellow “streaks” and looks different. He has a history of recurring tinea pedis.
- He has diabetes and is on metformin BID and a small dose of Humulin N at bedtime. He started swimming a year ago to improve his health after he had a “mild” heart attack.
- Upon examination, you notice a yellowish discoloration mainly under the distal end of a thickened toenail.

Risk factors for onychomycosis⁹

- Risk factors include: age (increased risk with older age), gender – males 2.4x at risk than females¹⁰, history of tinea pedis or known infected family members.
- Medical conditions that increase risk of infection include diabetes, immunodeficiency, psoriasis or genetic factors.
- Other contributory factors include: poor peripheral circulation, nail trauma, occlusive shoes, smoking, sports activities or other activities involving bare feet.

When to consider treatment

- Patients with diabetes and/or additional risk factors for cellulitis (i.e. prior cellulitis, venous insufficiency, edema). Onychomycosis may be a predictor of foot ulcer in a diabetic patient¹¹.
- Patient experiencing nail pain or discomfort.
- Cosmetic improvement desired.

Diagnosis

- Nail clippings, scrapings under the nail and deep nail samples are essential to confirm diagnosis of dermatophyte infection. This is recommended **before** starting treatment!
- If negative for dermatophytes, assess for possible psoriasis, lichen planus, nail trauma, onycholysis (e.g. distance runners), changes due to aging or gel nails, & yellow-nail syndrome.

Other Fungal Infections: Clinical Pearls from the Antifungal Chart (chart, next page &/or online)

Common skin infections

- Nystatin **only** effective for *Candida* infections (e.g. diaper rash, intertrigo, vulvovaginal infection).
- Combination products that contain steroids and/or nystatin should **not** be used for dermatophyte infections (e.g. Viaderm[®]: nystatin, neomycin, gramicidin & triamcinolone; Lotriderm: clotrimazole + betamethasone).

Oral treatment

- **Terbinafine LAMISIL 250mg PO once daily** is the drug of choice (cure rate >50-80%, however relapse is common). Terbinafine is more effective than itraconazole¹² and able to maintain cure for a longer duration (2 year follow-up).¹³ Terbinafine also has less risk for potential drug interactions.
- Alternate treatments
 - Itraconazole *SPORANOX* pulse therapy is an alternative if terbinafine contraindicated.
 - Fluconazole *DIFLUCAN* is less effective but is useful in patients unable to take the above.

Duration & approach to treatment^{14,15}

- Duration of treatment for terbinafine and itraconazole:
 - **toenail 12-16 weeks**; fingernail 6 weeks.
- **Weekly topical terbinafine cream** application after completion of oral treatment may be tried to prevent reinfection (expert opinion). The cream is applied between toes and around nail margin.
- Alternate treatments
 - Itraconazole pulse therapy (ie. 200mg po BID for 1 week per month) may decrease costs, side effects when compared to fixed dose (ie. 200mg po daily). Cure rates are similar with pulsed vs. continuous treatments. {Continuous daily dosing is more effective than pulse therapy for terbinafine.}¹⁶
 - Fluconazole 150mg po once weekly (x 6-12 months for toenail; x ≥3 months for fingernail).^{17,18}
- To monitor for treatment success, **mark the nail at completion of oral treatment**. This can be done by filing a line in the nail at the proximal part of known infection and marking with a permanent marker. Ask the patient to return if mark and affected toenail do not grow out or if infection moves proximal past the marked line.

Cautions including contraindications and side effects

- A meta-analysis¹⁹ found the risk of severe liver injury or asymptomatic elevations of serum transaminases with all treatments to be <2%. Liver enzymes should be done at baseline and after 4-6 weeks with terbinafine and monthly for itraconazole.
- Itraconazole is contraindicated in patients with heart failure or ventricular dysfunction and in patients using drugs metabolized by CYP 3A4 (see Antifungal Chart).

Oral candidiasis

- The nystatin dose for oral candidiasis (adult) is usually **5ml** QID to ensure enough liquid to cover area in mouth

Vulvovaginal candidiasis (uncomplicated)

- **1-3 days** with a topical azole as effective as 6-7 days for treatment but allow ~3 days for symptom resolution.
- 7 day topical azole treatment recommended in pregnancy

Select drug interactions with antifungals²⁰

- **Terbinafine** has minimal significant drug interactions and is a good antifungal option for patients on multiple drug regimens. As an inhibitor of CYP 2D6, it does still have some potential for drug interactions including increasing the levels and effect of TCAs, beta-blockers and antipsychotics. (See also Antifungal Treatment Chart.)
- **Itraconazole** is a **strong CYP 3A4 inhibitor** resulting in many frequent and significant drug interactions. The majority of drug interactions result in increased levels of drugs that may: prolong QT interval (i.e. amiodarone, quinidine, erythromycin), increase side effects (digoxin-nausea, vomiting; nifedipine-hypotension, dizziness; simvastatin/lovastatin-rhabdomyolysis; repaglinide, pioglitazone?-hypoglycemia) or increase toxicity (i.e. cyclosporine, tacrolimus)
 - Strong CYP 3A4 inducers (i.e. phenytoin, grapefruit juice) and antacids may decrease itraconazole levels.
- **Fluconazole** has less potential for major drug interactions than itraconazole because of its renal elimination and lesser effects as an enzyme inhibitor. (Agent is 3rd line in onychomycosis due to limited efficacy.)

Is ciclopirox nail lacquer ^{Penlac} an option?²¹

- Penetration into the nail is limited and use is of minimal value. It is slightly more effective when compared to placebo²²; no additive benefit when combined with oral terbinafine²³
- Recurrence is common on discontinuation.
- Consider cost of solution: \$140 / 12gm bottle
- The application process may be difficult for elderly & those with vision impairment. {Daily application 5mm beyond nail margin, on the bottom of the nail and skin under nail recommended. Remove **weekly** with isopropyl alcohol, trim or remove any damaged nail.} Treat x 48 weeks.

Home remedies – Do they work?

- Home remedies like vinegar, *Listerine*, *Vicks Vaporub*, vitamin E or thyme oil have no proven benefit.
- There is minimal evidence to support use of tea tree oil. It is a potent sensitizer and can cause local irritation and inflammation, producing skin reactions similar to those seen with poison ivy.²⁴



RxFiles Academic Detailing Team
out and about in SK



Best Educational Booth
FMF – Calgary – Oct 2010

References – RxFiles Newsletter : Antifungal newsletter (April 2010)

- 1 Roberts DT, Taylor WD, Taylor WD, Boyle J. Guidelines for treatment of onychomycosis. *Br J Dermatol* 2003;148:402-10
- 2 de Berker D. Clinical practice. Fungal nail disease. *N Engl J Med*. 2009 May 14;360(20):2108-16.
- 3 Crawford F, Hollis S. Topical treatments for fungal infections of the skin and nails of the foot. *Cochrane Database of Systematic Reviews* 2007, Issue 3. Art. No.: CD001434. DOI: 10.1002/14651858.CD001434.pub2
- 4 Medical letter. Treatment guidelines : Antifungal drugs. *The Medical Letter* 2008(Jan);6(65):1-8. (Medical Letter-Treatment Guidelines-Antifungal drugs. Dec. 2009)
- 5 Goldstein AO, Goldstein BG. Onychomycosis. Up-to-date. Accessed 18 August 2009
- 6 Olde Hartman TC, van Rijswijk E. Fungal Nail Infection. *BMJ* 2008 Jul 10; 337: a429
- 7 Hainer BL. Dermatophyte infections. *Am Fam Physician*. 2003;67:101-8
- 8 Seebacher C, Brasch J, Abeck D, et al. Onychomycosis. *Mycoses* 2007;50:321-7
- 9 Sigurgeirsson B, Steingrimsdottir O. Risk factors associated with onychomycosis. *J Eur Acad Dermatol Venereol* 2004 Jan;18(1):48-51
- 10 Gupta AK, Jain HC, Lynde CW, Macdonald P, Cooper EA, Summerbell RC. Prevalence and epidemiology of onychomycosis in patients visiting physicians' offices: a multicenter canadian survey of 15,000 patients. *J Am Acad Dermatol* 2000 Aug;43(2 Pt 1):244-8.
- 11 Boyko EJ, Ahroni JH, Cohen V, et al. Prediction of diabetic foot ulcer using commonly available clinical information: the Seattle Diabetic Foot Study. *Diabetes Care* 2006;29:1202-1207.
- 12 Crawford F, Young P, Godfrey C et al. Oral treatments for toenail onychomycosis. *Arch Dermatol* 2002;138:811-6.
- 13 De Cuyper C, Hindryckx PH. Long-term outcomes in the treatment of toenail onychomycosis. *Br J Dermatol* 1999 Nov;141 Suppl 56:15-20.
- 14 Sigurgeirsson B, Olafsson JH, Steinsson JB, et al. Long-term effectiveness of treatment of treatment of terbinafine vs itraconazole in onychomycosis: a 5-year blinded prospective follow-up study. *Arch Dermatol* 2002;138:353-7.
- 15 Evans EG, Sigurgeirsson B. Double blind, randomized study of continuous terbinafine compared with intermittent itraconazole in treatment of toenail onychomycosis. *BMJ* 1999;318:1031-35
- 16 Warshaw EM, Fett DD, Bloomfield HE, Grill JP, Nelson DB, Quintero V, Carver SM, Zielke GR, Lederle FA. Pulse versus continuous terbinafine for onychomycosis: a randomized, double-blind, controlled trial. *J Am Acad Dermatol* 2005 Oct;53(4):578-84.

Case Discussion (continued):

- Nail clipping and scraping was cultured and came back positive after 4 weeks. Due to patient's diabetes, potential risk for cellulitis and history of tinea pedis, it was decided to recommend pharmacological treatment.
- The option of treating, including the benefits, risks and costs were discussed. Since he had diabetes, he was deemed to derive substantial benefit.
- Terbinafine 250mg once daily x 12 weeks was initiated
- Mr T. returned 3 months later after completing a course of treatment and noticed an improvement in his toe appearance. However, it still did not look "normal". He was reassured that he did not require additional treatment at this time. The nail was marked at the margin proximal to the infection and patient counseled to return if the infection moved past the mark or failed to grow out in the coming 12-18 months. He was instructed to trim & file the nail as it grew.

Prevention topics to discuss with patient...

- *Treatment of tinea pedis*
- *Proper footwear e.g. wear sandals/slippers in communal areas such as swimming pools, locker rooms, gyms, mosque, etc.*
- *Avoid going barefoot where possible*
- *Proper nail hygiene – trim nails short & straight across*
- *Avoid using same nail clippers or files on both diseased and normal nails; have separate tools for infected nails or disinfect between use*
- *Disinfection of socks & shoes*
- *Clean bathroom surfaces with bleach*

Coming soon ...

- ◆ **Summer 2010: RxFiles Drug Comparison Charts book – 8th Ed.**
 - è ~140 pages; 14 new charts (e.g. anti-infectives for common infections, CKD, osteoporosis, sexual dysfx, SMBG, substance abuse, transplantation drug tx considerations, vaccines (adult), etc
 - è Pre-release ordering now available. See our online store or form: <http://www.rxfiles.ca/rxfiles/uploads/documents/1A-CHT-Book-ORDERFORM.pdf>
- ◆ **Information Mastery Course – Saskatoon, May 7-8, 2010**
 - è a practical approach to evidence based medicine for clinicians
 - è guest faculty from Tufts School of Medicine/Health Care Institute
 - è limited registration space for this very special event
 - è co-hosted with Continuing Professional Learning, U of S. <http://www.rxfiles.ca/rxfiles/uploads/documents/Information-Mastery-Course.pdf>

Acknowledgements: Dr. Hull (SHR-Dermatology); Dr. Lichtenwald, (Dermatology), Dr. Yvonne Shevchuk (College of Pharmacy, U of S); Dr. Sanche (SHR-Infectious Disease), Dr. T. Laubscher CCFP (FM, U of S), M. Jin (Pharm D, Hamilton), A. Bhalla (Pharm D, Ontario)& the RxFiles Advisory Committee *Shannon Stone BSP, Loren Regier BSP, BA, Brent Jensen BSP*

DISCLAIMER: The content of this newsletter represents the research, experience and opinions of the authors and not those of the Board or Administration of Saskatchewan Health Region (SHR). Neither the authors nor Saskatchewan Health Region nor any other party who has been involved in the preparation or publication of this work warrants or represents that the information contained herein is accurate or complete, and they are not responsible for any errors or omissions or for the result obtained from the use of such information. Any use of the newsletter will imply acknowledgment of this disclaimer and release any responsibility of SHR, its employees, servants or agents. Readers are encouraged to confirm the information contained herein with other sources. Additional information and references online at: www.RxFiles.ca

- 17 Scher RK, Breneman D Rich P, et al. Once-weekly fluconazole (150,300, or 450mg) in the treatment of distal subungual onychomycosis of the toenail. *J Am Acad Dermatol* 1998;38:S77-86.
- 18 Brown SJ. Efficacy of fluconazole for the treatment of onychomycosis. *Ann Pharmacother* 2009;43:1684-91
- 19 Chang CH, Young-Xu Y, Kurth T, et al. The safety of oral antifungal treatments for superficial dermatophytosis and onychomycosis: a meta-analysis. *Am J Med* 2007;120(9):791-8.
- 20 Bruggemann RJ, Alfenaar JW, Bijllevens NM, et al. Clinical relevance of the pharmacokinetic interactions of azole antifungal drugs with other coadministered agents. *Clin Infect Dis*. 2009 May 15;48(10):1441-58.
- 21 Hart R, Bell-Syer SE, Crawford F, Torgerson DJ, Young P, Russell I. Systematic review of topical treatments for fungal infections of the skin and nails of the feet. *BMJ*. 1999 Jul 10;319(7202):79-82
- 22 Crawford F, Hollis S. Topical treatments for fungal infections of the skin skin and nails of the foot. *Cochrane Database Syst Rev* 2007; (3) CD001434.
- 23 Gupta AK. Ciclopirox topical solution, 8% combined with oral terbinafine to treat onychomycosis: a randomized, evaluator-blinded study. *J Drugs Dermatol* 2005;4:481-485
- 24 Natural Medicine Comprehensive Database. Tea tree oil monograph. Accessed 2 October, 2009. (<http://www.naturaldatabase.com/>)

Pages 1 & 2 of the Antifungal Drug Comparison Chart are included with this newsletter. These pages include the antifungals most used in primary care. Go online to www.RxFiles.ca where the complete antifungal drug chart can be found which has a 3rd page covering several other antifungals (e.g. ketoconazole **NIZORAL**, voriconazole **Vfend**, posaconazole **POSANOL**, caspofungin **CANCIDAS**, micafungin **MYCAMINE**, anidulafungin **ERAXIS**, & amphotericin-B **FUNGIZONE**, **ABELCET**, **AMBISONE**).

Produced by RxFiles – a provincial academic detailing service funded by Saskatchewan Health. For more information check our website at www.RxFiles.ca or contact us c/o Saskatoon City Hospital, 701 Queen Street, Saskatoon, SK. S7K 0M7 Phone (306) 655-8505.

Copyright 2010 – RxFiles, Saskatoon Health Region (SHR) www.RxFiles.ca



Onychomycosis 8,9

Key signs: nail thickening, discoloration, & separation from nail bed.^{10,11,12,13} **C&S to confirm** prior to tx. (Clip, scrape & deep nail sample to avoid false negatives.) **Cause:** toenail→commonly dermatophytes; fingernail→may be yeast¹⁴ [yeast e.g. candida; dermatophyte=filamentous fungi (eg. tinea)]
Pearls: uncommon to have finger without toenail involvement; file & mark margin of fungus on nail at completion of tx to monitor success!
Risk factors: ↑ prevalence with ↑ age (15-20% in pts ≥ 40 yrs); swimming, barefoot, tinea pedis, diabetes, immunodeficiency, living with an infected family member^{15,16}
Tx: □ **Oral terbinafine** or itraconazole: **x12-16wks** toe; success:50-80%; relapse: ~25-30%¹⁷; topical terbinafine weekly to prevent relapse? {Effectiveness: terbinafine > itraconazole >> fluconazole if unable to tolerate other tx; consider cost, success rate, SE risk}¹⁸
 □ Itraconazole pulse tx less \$\$ & SE, but requires scheduling; however terbinafine pulse treatment lower cure rate than daily dose¹⁹
 □ **Topical:** **Nail lacquer** in mild, distal dx, minimal penetration; combo with po **no** added benefit
Prevention: tx tinea pedis; wear sandals/slippers in communal areas bathing places, locker rooms, gyms, mosque
 □ **Home remedies** eg. Vicks VapoRub, vinegar no proven tx benefit. Tea tree oil: little evidence for benefit²⁰; allergy.

Oral Candidiasis 21,22

Key Signs: Pseudomembranous form: white plaques on oral mucosa; atrophic form: erythema without plaque (common in elderly with dentures denture stomatitis). Angular cheilitis may be present.
Causes: commonly *Candida albicans* **Risk factors:** smoking, poor dental hygiene, inhaled or systemic **steroid use**, antibiotics, diabetes, immunodeficiency, ↓ saliva
Tx: □ **Mild dx:** Topical nystatin or oral fluconazole effective x **7days** minimum (or 2+days after improved.)
 □ **Dentures:** **disinfect** chlorhexidine rinse ~20-30min & tx with topical antifungal to mucosa & denture base²³.
 □ **Refractory, recurrent or esophageal** infections need systemic azoles fluconazole; topical tx ineffective. May indicate compromised immune system; consider referral to ID (? HIV).
Prevention: If on **inhaled steroid**, use aerochamber, rinse mouth & spit after each use.
Dentures: daily cleaning recommended (chlorhexidine useful, rinse well)²⁴; +/- nystatin but not at same time
Infant: □ Nystatin safe, ↓cost but ↓effective → poor oral adherence & QID. comparison data limited^{25,26}
 □ Fluconazole more effective, once daily dosing but ↑ cost; **not** officially approved in newborns.
 □ Gentian violet 0.5-1% aqueous soln BID effective, but longer tx period, messy, & associated with ulceration.^{27,28}
 □ Breastfeeding infant: consider topical tx of nipple²⁹ (eg. clotrimazole, miconazole, nystatin) lack safety data

Common Fungal Skin Infections 36,37,38











Causes: Candida, epidermophyton, trichophyton, microsporium **Risk factors:** animal exposure (ie. vets, vet techs), skin trauma (ie. wrestlers), diabetes, immunodeficiency, ↓ circulation, poor hygiene, warm/humid climate.
General tx info: Apply antifungal to **affected & surrounding area (1-2 inches beyond rash)**.
 □ Continue x **1wk** after sx's gone & skin looks healed to ensure eradication (often ~10-14 days).
 □ Keep area clean & dry (use non-scented talc or powder baby powder, Goldbond, tolnaftate as prophylaxis).
 □ Nystatin **not useful** for dermatophyte infections; effective for candidal infections.
 □ **Oral tx:** nail, scalp Kerion: inflamed purulent mass, from livestock? add prednisone, beard, severe/widespread or if recurrent.
 □ Combination with **steroids not usually recommended** due to ↑ SE, cost & ↓ cure rates.
Prevention: Avoid sharing personal items & towels. Avoid wearing tight or occlusive clothing.
 Wash linens & clothing in hot water & hot dryer or line dry & expose to UV rays; disinfect shoes.
 i) **Seborrheic dermatitis:**³⁰ Commensal overgrowth of yeast. Topical/shampoo azoles & ciclopirox olamine useful. Intermittent shampoo use once weekly or every other week after tx may ↑ remission. (limited comparison data)
 ii) **Tinea capitis (Scalp):** **Common in kids** cats, cows; oral terbinafine_{DOC} x **4-8wks** +/- selenium sulfide shampoo 2-3x per wk (x5mins) to ↓ spread. Other options: oral fluconazole, itraconazole, (griseofulvin).
 iii) **Tinea corporis (Body):** Tx options: topical azoles (clotrimazole, miconazole) & terbinafine. Consider topical azoles first, terbinafine **slightly** more effective/rapid but ↑ cost. Tx: x**2-4** wks.
 iv) **Tinea Cruris (Groin):** Common in adolescent & young adult ♂; if wear tight jeans/pantyhose. Overdiagnosed?
Tx: Topical azole clotrimazole, miconazole x 2-4wk or terbinafine cream/spray daily x 2-4wk. Assess for tinea pedis.
 v) **Tinea pedis (Foot):** **Tx** Effective: terbinafine > azole (clotrimazole, miconazole) > tolnaftate; consider cost & dosing schedule³¹. **Treat topically x 4wks.** (Common: elderly dry cracked skin; adolescent between toes.)
 vi) **Tinea Pityriasis versicolor:**^{32,33} Commensal overgrowth of Malassezia yeast. Use topical antifungals 1st mild dx. Apply azole to whole affected area (ie. chest) every day x 1wk, then q. weekly for prophylaxis). If severe/recurrent consider **short-term** 1-5 days po (keto-, flu-, itra-conazole (↑ SE)). **Oral terbinafine ineffective**³⁴. Suggest selenium sulfide 2.5% or ketoconazole 2% shampoo ↓ recurrence weekly or 1-2x /month x 40+ yrs (ie. long-term)
Candidal Intertrigo³⁵: Common in moist skin folds (especially in obese, ostomy, etc.); results in tender, burning, pruritic areas with satellite lesions; **Tx:** consider nystatin **powder**, topical antifungals

Vulvovaginal Candidiasis 39

Key signs: pruritus, soreness, dyspareunia, external dysuria; possibly thick & curdy discharge
Causes: *Candida albicans*, occasionally non-albicans; associated with antibiotic use; **rule out UTI/STI**
Tx: Topical azoles (see table) or oral fluconazole. Oral route often preferred by pts; consider cost.⁴⁰
 {Cochrane: no difference in effectiveness of fluconazole oral vs intra-vaginal OTC routes}
 □ 1-3days topical as effective as 6-7days with better compliance. **Allow ~3 days for sx resolution.**
 □ **Recurrent** cases (≥4/yr) may benefit by addressing risk factors uncontrolled diabetes, high dose estrogen OC (?HIV); try 1) longer initial course of topical (7-14days) then clotrimazole 200mg pv 2x weekly or 500mg Supp pv weekly; or 2) fluconazole 150mg q72h x 3 doses⁴¹ then fluconazole 150mg po weekly. Treat male partner?: controversial, but may benefit if *Candida balanitis* present.; tx-topical azole BID x 1 week^{42,43,44}
 □ **Complicated** vaginitis...10% : ≥7days topical tx or fluconazole 150mg q72hr for 3 doses-IDSA guidelines
 □ **Pregnancy:** requires longer tx interval (eg. 7 days azoles; 14 days nystatin; 1 day fluconazole po) topical azole (clotrimazole, miconazole) more effective & convenient than nystatin; tx topical 1st line systemic absorption low; ↓ risk of birth defects⁴⁵; oral fluconazole 2nd line Avoid 1st trimester & ≥ 400mg daily as teratogenic.
 □ Topical boric acid 600mg cap PV hs x2wks an option if *C. glabrata* (rare); compounded not commercially available⁴⁶
 □ **Dietary yogurt (with live culture)** or oral Lactobacilli caps: do **NOT** prevent post-antibiotic vulvovaginitis, but may help restore normal flora⁴⁷ {Vaginal yogurt controversial.}
 □ topical vaginal tx containing mineral or vegetable oil {e.g. miconazole vaginal ovules problem} may ↓ effectiveness of **condoms**, or other vaginal contraceptive devices (eg. **diaphragms**) during treatment & up to 3 days post-tx⁴⁸ {Okay: clotrimazole products & miconazole cream.}
 □ **Apply antifungal underneath barrier cream** until rash is resolved.
 □ Topical nystatin, clotrimazole, miconazole, or ketoconazole if rash candidal or >3 days.
 □ **Combo topical corticosteroid/antifungal** products **not** routinely recommended as may result in dilution, ↑ SE & mask Sx of infection. eg. Viaderm-KC, Kenacomb If necessary: use only **low-potency, short-term corticosteroid!!!** **Best to apply creams separately** allowing a few minutes between applications. {Alternately, add hydrocortisone powder 1% to azole cream. See also OTC dermatology section.}

Diaper Rash 49

Antifungals: Topicals & Vaginal:	Tinea pedis/cruris/corporis	Comments:	Vaginal candidiasis All OTC	Cost
<p>Ciclopirox olamine LOPROX 1% top crm (45gm); 1% top lotion (60ml) PENLAC 8% Nail lacquer X; STIEPROX 1.5% Shampoo X (100ml)</p> <p>Clotrimazole CANESTEN 1% top crm (15,30 & 500gm); 200,500mg vag tab; 1, 2 & 10% vag crm Generic OTC 1% top crm (20,30,50 & 500gm); 1, 2% vag cream [higher % for shorter term tx]</p> <p>Ketoconazole Generic 2% top crm (30gm) NIZORAL 2% Shampoo X (60,120ml)</p> <p>Miconazole MONISTAT-DERM 2% top crm (15,30gm) MONISTAT, Generic 100, 400, 1200mg vag ovules; 2, 4% vag cream;</p> <p>Nystatin MYCOSTATIN, Generic 100,000 U/G top crm & oint (bulk powder available for compounding topical powder) OTC (15,30 & 450gm); 25,000 & 100,000 U/G vag cream</p> <p>Terbinafine LAMISIL 1% crm (15,30gm); 1% top spray soln (30ml)</p> <p>Tolnaftate TINACTIN 1% top crm; powder; soln; top spray Others(Undecylenic acid-Desenex / Fungicure, Tolnaftate-Dr. Scholl's OTC products): less data, less effective</p>	<p>Apply bid x 2-4 weeks</p> <p>Apply bid x 2-4 weeks</p> <p>Apply once daily x 2-4 wk (x 6wks tinea pedis)</p> <p>Apply bid x 2-4 weeks</p> <p>Nystatin NOT effective for dermatophytes!</p> <p>Apply daily x 2-4wk (x 1-2wk mild tinea pedis)</p> <p>Apply bid x 2-4wks</p>	<p>□ Cost Considerations: - terbinafine more expensive but more rapid effect ∴ azoles generally used first; consider amount of product required, dosing schedule & length of tx - Cost/30gm tube: clotrimazole \$12-15; miconazole \$12-15; terbinafine \$20-25</p> <p>□ Consider oral tx if widespread, recurrent or failure with topical tx</p> <p>□ Creams or spray soln preferred over powders, except in skin folds.</p>	<p>CANESTEN 1 Combi Pak 500mg pv / 1%crum ▼ or Cream 10% x 1 day, CANESTEN 3 Combi Pak 200mg pv / 1%crum ▼ or Cream 2% x 3 days, CANESTEN 6 Cream 1% x 6 days. MONISTAT 1 Vag Ovule 1200mg x 1 day or Combi Pak 1200mg/2%crum x 1 day, MONISTAT 3 Dual Pak 400mg pv / 2%crum ▼ or Vag Ovule 400mg ▼ or Vag Cream 4% x 3 day, MONISTAT 7 Dual Pak 100mg pv / 2%crum ▼ or Vag crm 2% x 7day.</p> <p>TERAZOL 3 Supp 80mg x ▼ or Dual Pak 80mg pv / 0.8% crm ▼ or Vag crm 0.8% x 3day TERAZOL 7 0.4% crm ▼ x 7 day. CanesOral fluconazole 150mg po; & CombiPAK</p>	<p>\$14-18</p> <p>\$16-20</p> <p>\$20-30</p> <p>\$25-33</p>

Generic/TRADE (Strength & forms) g=generic	P 50	Side effects / Contraindications C Cautions	√ = therapeutic use / Comments / Drug Interactions DI (not exhaustive) ⁵¹ / Monitor M	INITIAL; MAX /USUAL DOSE {Drug of Choice highlighted in brown.}	\$  /course
Terbinafine HCL ▼ g Lamisil 250mg tab ⁵	B	Common: PO: headache, GI diarrhea, dyspepsia, abdominal pain, taste disturbance may persist after tx stopped, rash mild Serious: (≥0.01% to 0.1%) ↑AST & ALT or hepatotoxicity, (≤0.01%) SJS, toxic epidermal necrosis, erythema multiforme, pancytopenia, neutropenia Precaution: liver/kidney disease, lupus erythematosus	√ Onychomycosis & skin infections due to dermatophytes Tx severe tinea corporis, cruris, pedis unresponsive to topicals DI: CYP2D6 inhibitor: ↑effect of: TCA ↑TCA level, Possible: Beta blockers & Antipsychotics M: LFT's at baseline & at 4-6 wks of tx ⁵²	Onychomycosis: 250mg po daily (Fingernail: x 6wks; Toenail: x12-16 wks) Tinea capitis: 250mg po once daily x 4-8wk Pediatric dosing ≥ 4 yrs: (e.g. Tinea capitis x4wk) <20kg: 62.5mg/day po, 20-40kg: 125mg/day po, >40kg: 250mg/day po ⁵³	108/6wks 225/12wk 41-75/ 2-4wks
Fluconazole g Diflucan (50, 100mg tab) ▼   ; 150mg cap ▼, regular benefit SK formulary [CanesOral: new OTC formulation of fluconazole 150mg tab +/- clotrimazole 1% vag cream] 10mg/ml powder for oral suspension (P.O.S.) Diflucan IV soln 200mg/100ml vial, 400mg/200ml vial	C	Common: well tolerated; headaches, GI upset, rash Serious: Stevens-Johnson syndrome(SJS), hepatotoxicity, QT prolongation C: cisapride: ↑↑ drug level cause ↑QT & torsades des pointes; ergot alkaloids : ↑↑ ergot levels Cautions: -High dose ≥ 400mg/d in pregnancy & 1st trimester. -Pts on rifampin, phenytoin, valproic acid, isoniazid & po sulfonyleureas may be at ↑ hepatic risk. Thrush in Newborns: NOT officially indicated but is an off-label, more effective alternative to nystatin. - Full-term (37-44 wk GA) & 0-14 days: 3mg/kg q48h - Full-term (37-44wk GA) & >14 days: 3mg/kg q24h ⁵⁴ Dose varies on site &/or severity of infection	√ Active against most <i>Candida</i> species except <i>C.krusei</i> & some <i>C. glabrata</i> , <i>Coccidioides</i> , <i>Histoplasma</i> , <i>Cryptococcus</i> sp. in high doses Consider for oropharyngeal, esophageal or vaginal candidiasis DI: ↓ fluconazole level: rifampin. [Less DI's than azoles in general.] Moderate CYP3A4 inhibitor: ↑level of alfentanil, carbamazepine, cyclosporine, midazolam, quinidine, rifabutin, statins, tacrolimus, & triazolam. Strong CYP 2C9,2C19 inhibitor: ↑level of ergot alkaloid, glibenclamide, nevirapine, phenytoin, warfarin, zidovudine. Prolong QT interval: amiodarone, cisapride, clarithromycin, TCA's Renal dx: no adjustment needed for single-dose vaginal candidiasis M: liver enzymes, renal function; baseline & periodically if risk factors/long-term tx Comments:  Bioavailability of PO similar to IV; use PO if possible  ↓ DI due to ↑ renal excretion ~80% & ↓ hepatic metabolism effect  Compatible with breastfeeding  May require dose ↑ if obese with severe/systemic infection	Dose range: 100-800mg /day. Pediatric: 3mg/kg/day-12mg/kg/day. {5 adult dose.} Onychomycosis: 150mg po once weekly (Fingernail: x 3mos; Toenail: x 6-12mos) ⁵⁵ (3rd line adults; useful if ++DI's, peds pts) Oropharyngeal candidiasis: Load: 200mg po x1 →100mg po daily x 7 day (Peds: Load 6mg/kg → 3mg/kg/day x 14day) Esophageal candidiasis: 200-400mg od x 2-3wk Tinea versicolor: 400mg po x 1 dose Vulvovaginitis candidal: 150mg po once ^{OTC} Candidemia neutropenic & non-neutropenic: Load day 1:800mg →400mg daily until 14day post-signs/sx & after last +ve blood culture ; obese patients: consider 6-12mg/kg IDSA ⁽⁵⁶⁾	141/3mos 282/6mos 64 /wk 178-349 /2 wks 32 17 178/wk
Itraconazole ▼  Sporanox 100mg cap [Give cap with food acidic PH ↑ absorption; In past, was often given with cola.] 10mg/ml solution -soln more bioavailable than cap ⁵⁷ ; solution preferred for oral/esophageal candidiasis. [Take on empty stomach] **Dosage forms NOT interchangeable**	C	Common: dose-related nausea, diarrhea, abdominal discomfort, rash, edema, hypokalemia, ↑ transaminases, & dizziness Serious: SJS, hepatotoxicity failure, HF dose related negative inotropic effect at 400mg/d C: pts with ventricular dysfunction or HF; pts on negative inotropics or erythromycin; pts using drugs metabolized by CYP 3A4 (ie. cisapride, dofetilide, eletriptan, ergot alkaloids, lovastatin, midazolam, nisoldipine, pimozide, quinidine, simvastatin, triazolam); : pregnant women Caution: hepatic dysfunction, pts at risk for arrhythmias [See note at bottom for "Hepatic Risk" comment.]	√ Broader spectrum of activity than fluconazole: including <i>Candida</i> spp., <i>Cryptococcus neoformans</i> , <i>Aspergillus</i> spp., <i>Blastomyces dermatitidis</i> , <i>Coccidioides immitis</i> , <i>Histoplasma capsulatum</i> & dermatophytes. Consider for fluconazole resistant mucosal candidiasis DI: Strong CYP3A4 inhibitor: ↑ level of: amio-/drone-darone, astemizole, atorvastatin some, buspirone, CCB nifedipine, nisoldipine, felodipine, cisapride, cyclosporine, digoxin, dofetilide, eletriptan, ergot alkaloids, fentanyl, indinavir, lovastatin, midazolam, pimozide, quinidine, ritonavir, saquinavir, simvastatin, sirolimus, steroids ↑ level: budesonide, dexamethasone, flucicasona, methylprednisolone, tacrolimus, triazolam & vincristine. ↑ itraconazole level: indinavir, ritonavir ↓ itraconazole level: antacids, H2 receptor blockers, PPI due to ↓ acidity; carbamazepine, efavirenz, grapefruit juice, nevirapine, phenytoin, rifampin, rifabutin ↓ levels of oral contraceptives. ↑ level of: warfarin M: liver enzymes (every month if on long-term tx ie >1month) Comments:  most DI's, ↑ toxicity compared to other azoles	Dose range: 100-400mg/day Onychomycosis (if terbinafine contraindicated) Toenail: 200mg po daily x12wks or "pulse" tx: 200mg po BID x 1wk (3wks off & rpt 1wk x 2 cycles) Fingernail: 200mg po daily x 6wks or "pulse" tx: 200mg BID x 1 wk (3wks off & rpt x 1 wk) Oropharyngeal candidiasis: if fluconazole resistant 200mg po once daily of soln x 14 days Esophageal candidiasis: if fluconazole-resistant 200mg po daily of soln x 14-21 days Tinea versicolor: 200mg po daily x 5-7 days (pyriariss versicolor) or 400mg x 1 dose ^{58,59} Caps less expensive (~half the cost) but less bioavailable; solution used for pricing of oral/esophageal candidiasis only.	822 /12wks 408/6wks (daily dose) 423/3mos 282/2mos (pulse tx) 283/ 14days
Nystatin ▼ g 500,000 unit tab 100,000 units/ml susp	A C	Common: well-tolerated; nausea, vomiting, diarrhea at high doses Caution: contains sucrose; may ↑ risk for dental caries	√ Fungi-static & cidal; may be used for candidal skin infections, Oropharyngeal & vulvovaginal candidiasis; for topical skin & vaginal candidal infections during pregnancy  slightly less effective for most conditions but safe, inexpensive	Children & adults: {liquid; swish & swallow!} Thrush (mild): 500,000units (5ml) qid x 7days or 2days after improvement. Pediatric: [may use 0.5ml & swab for infants] Infants ♂ thrush: 100,000-200,000 units qid	15 / 7days

↓ dose for renal dysfunction ζ =scored tab χ =Non-formulary SK \equiv =Exception Drug Status SK \otimes =not covered by NIHB \blacktriangledown =covered by NIHB \varnothing =prior NIHB approval CCB=calcium channel blocker **C**=contraindication crm=cream DI=drug interaction DOC=drug of choice Dx=disease fx=function g=generic avail. GA=gestational age GI=gastrointestinal HF=heart failure LFT=liver function tests n/v=nausea/vomiting OC=oral contraceptive OTC=over the counter pc=after meals po=oral PPI=proton pump inhibitor **Pr**=prescription Pt=patient pv=per vagina SAP=special access program SE=side effect SJS=Stevens-Johnson syndrome STI=sexually transmitted infection Sx=symptoms TCA=tricyclic antidepressant Tx=treatment UTI=urinary tract infection vag=vaginal wt=weight

When choosing drug keep in mind: frequency of dosing, dosing with regards to food, & organism coverage.

Comments: When not to use fluconazole: positive fungal urine cultures without symptoms of upper genitourinary disease, systemic candidiasis, or an impending genitourinary tract procedure; positive sputum cultures.
Special Considerations: Hepatic Risk: Overall incidence <2% for all; for oral tx of onychomycosis treatment: ketoconazole>itraconazole>terbinafine. Pulse treatment may reduce risk, but less effective for terbinafine.
Useful links: www.dermnet.com www.RxFiles.ca See page 53 (book or online) for: voriconazole ^{VFEND}, posaconazole ^{SPRIAFIL, POSANOL}, ketoconazole, echinocandins ^{CANDIDAS, MYCAMINE, ERAXIS}, amphotericin B. ^{FUNGIZONE, ABELCET, AMBISOME}
Other drugs: flucytosine ^{SAP} – add-on po tx of Candida endocarditis/meningitis with Amphotericin B. ^{griseofulvin} ^{FULVICIN}: not indicated in Canada but bulk supply available for compounding; is available in some areas of the world; especially useful in T. capitis; newer options available for tinea infection. ^{butoconazole} – 2% vag crm available, more expensive, no advantages over other indicated treatment for vaginal candidiasis; contains mineral oil: caution with condoms, diaphragms.
Investigational Drugs: Ravuconazole, Isavuconazole ^{invasive aspergillosis & candidiasis}, Pramiconazole & Albaconazole ^{onychomycosis}.

Acknowledgements: Contributors & Reviewers: Dr. P. Hull (MD, Dermatology, Saskatoon) Dr. D. Lichtenwald (MD, Dermatology, Saskatoon); Y. Shevchuk (PharmD, C of Pharmacy, U of S, Saskatoon); S. Sanche (MD, Infectious Diseases - Internal Med, Saskatoon), S. Skinner (MD, Infectious Diseases - Internal Med, Saskatoon), B.Tan (MD, SHR-Ped ID), M Jim (Pharm D, Hamilton), A Bhalla (Pharm D, Ontario) & the RxFiles Advisory Committee. Prepared by: Shannon Stone BSc, Brent Jensen BSc, Loren Regier BSc BA
DISCLAIMER: The content of this newsletter represents the research, experience and opinions of the authors and not those of the Board or Administration of Saskatchewan Health Region (SHR). Neither the authors nor Saskatchewan Health Region nor any other party who has been involved in the preparation or publication of this work warrants or represents that the information contained herein is accurate or complete, and they are not responsible for any errors or omissions or for the result obtained from the use of such information. Any use of the newsletter will imply acknowledgment of this disclaimer and release any responsibility of SHR, its employees, servants or agents. Readers are encouraged to confirm the information contained herein with other sources. Additional information and references online at www.RxFiles.ca
Copyright 2010 – RxFiles, Saskatoon Health Region (SHR) www.RxFiles.ca



<p>Ketoconazole ▼ </p> <p>Nizoral 200mg tab</p> <p>(see topicals section above for topical, shampoo)</p>	<p>C Common: poorly tolerated; anorexia, nausea, vomiting ^{high doses}; pruritus, rash dizziness, ↓ testosterone level: gynecomastia, ↓ libido & loss of potency in ♂, menstrual irregularities in ♀</p> <p>Serious: ↓steroidogenesis adrenal & ↓cortisol; hepatotoxic</p> <p>C: astemizole, cisapride, triazolam</p>	<p>√ Rarely used orally</p> <p>D: similar to itraconazole (see above)</p> <p>Strong CYP3A4 inhibitor: ↑ level of amio-drone-darone, cyclosporine, digoxin ^{potential}, ergot alkaloid, lovastatin, pimoizide, quinidine, rifabutin, simvastatin, tacrolimus, (similar to itraconazole)</p> <p>M: liver transaminases</p> <p>Comment: ☑With food & at bedtime to ↓SE ☑breastfeeding compatible</p>	<p>200; 400mg 200-400mg once daily at bedtime</p> <p>Pediatrics ≥ 2 yrs: 3.3-6.6mg/kg/day po once daily</p> <p>Tinea versicolor ^{60,61}: 400mg x 1 dose or (pityriasis versicolor) 200mg daily x 5-7 days</p>	<p>10 /400mg dose 15-17/ 5days</p>
<p>Voriconazole </p> <p>Vfend</p> <p>50, 200mg tab; (Good oral absorption)⁶² (Take on empty stomach)</p> <p>IV 200mg/vial</p> <p>Relatively new drug; often requiring Infectious Disease Service consult!</p>	<p>D Common: rash ~7%, photosensitivity, confusion, hallucinations, ↑ transaminases, transient visual disturbances ~20-23% including blurred vision, photophobia, & altered perception of color/image ^{may resolve early}</p> <p>Serious: SJS ^{rare}, hepatotoxicity</p> <p>C: astemizole, barbiturates, carbamazepine, cisapride, efavirenz, ergot alkaloids, pimoizide, quinidine, rifabutin, rifampin, high dose ritonavir >400mg BID, sirolimus, St. John's wort & terfenadine.</p> <p>: pregnant women</p> <p>Caution: hepatic dysfunction, pts at risk for arrhythmias</p>	<p>√ Similar spectrum to itraconazole;</p> <p>More active: <i>Aspergillus</i> spp & <i>Candida glabrata</i> & <i>krusei</i>, <i>Fusarium</i></p> <p>D: ↓ levels of voriconazole: barbiturates, carbamazepine, efavirenz, phenobarbital, phenytoin, rifampin, rifabutin, ritonavir, & St John's wort.</p> <p>Moderate CYP3A4 inhibitor ↑ levels of: alfentanil, amio-drone-darone, cisapride, cyclosporine, efavirenz, methadone, midazolam ^{po (& higher iv dose)}, omeprazole, sirolimus, tacrolimus, triazolam & vincristine</p> <p>Strong CYP 2C9, weak 2C19 inhibitor ↑ levels of: methadone, warfarin</p> <p>Liver dx: Initial loading dose, but half maintenance dose if liver cirrhotic</p> <p>Renal dx: if CrCl<50 ml/min-use only po formulation ^{solubilizing agent can accumulate}</p> <p>M: liver enzymes; serum level monitoring for serious infections only</p> <p>Comments: ☑ DOC-invasive aspergillosis ⁶³ ☑ serum levels may vary ⁶⁴ ☑ poor CYP2C19 metabolizers (ie Asian ~20-30%) ⁶⁵ ↑drug level</p>	<p>Dose range: 200-600mg/day</p> <p>Aspergillosis: 6mg/kg q12h x 1 day → then 4mg/kg or: if >40kg ♂ 200-300mg po q12h If <40kg ♂ 100-150mg po q12h Adjust dose based on levels if not responding. {Above dosing higher than previously recommended (200mg po q12h >40kg)}</p> <p>Oropharyngeal: if fluconazole resistant 200mg po bid x 14-21 day</p> <p>Esophageal candidiasis: if fluconazole-resistant 200mg po bid x 14-21 day</p>	<p>148 /200mg vial 1,509-2,259 /14-21 days</p>
<p>Posaconazole ⁶⁶ </p> <p>Posanol <i>Spirafil</i></p> <p>40mg/ml suspension (cherry flavored) (Take with high-fat meal or meal replacement to ↑ absorption)</p> <p>Relatively new drug; often requiring Infectious Disease Service consult!</p>	<p>C Common: fairly well-tolerated; diarrhea, nausea ~6%, vomiting, headache ~6%, hypokalemia ↑transaminases ^{similar to fluconazole}</p> <p>Serious: hepatic necrosis; QT prolongation & arrhythmias</p> <p>C: ↑level of astemizole, cisapride, ergot alkaloid, pimoizide, quinidine, sirolimus, terfenadine</p> <p>: pregnant women</p> <p>Caution: hepatic dysfunction, pts at risk for arrhythmias</p>	<p>√ Similar spectrum to itraconazole with activity against Zygomycetes (alternative to amphotericin B), <i>Cryptococcus</i>, <i>Aspergillus</i>; refractory oropharyngeal/esophageal candidiasis; prophylaxis of <i>Aspergillus</i> & <i>Candida</i> infection in neutropenics & stem cell transplant recipients; option for prophylaxis & tx of invasive fungal dx (broad spectrum; potentially less resistance)</p> <p>D: Moderate-strong CYP3A4 inhibitor ⁶⁷: ↑level of amio-drone-darone ^{theoretical}, atazanavir, cyclosporine, digoxin ^{potential}, midazolam⁶⁸, rifabutin, sirolimus, tacrolimus, terfenadine, triazolam & vincristine</p> <p>↓ levels of posaconazole: cimetidine, efavirenz, phenytoin, rifabutin.</p> <p>M: liver enzymes; electrolytes (K⁺, Mg⁺⁺, Ca⁺⁺)</p> <p>Comments: ☑ Less DI's; metabolized by glucuronidation</p>	<p>Consult with Infectious Disease Specialist/Service for Posaconazole use!</p> <p>Dose range: 100-800mg/day {Pts > 13yrs}</p> <p>Oropharyngeal candidiasis: Load: Day 1: 100mg bid → 100mg od x 13day</p> <p>Fluconazole-refractory oropharyngeal dx: 400mg po BID x3d → 400mg daily x 4wk ^{IDA} ⁶⁹</p> <p>Esophageal, fluconazole refractory: 400mg po BID x 14-21 day;</p> <p>Prophylaxis of invasive infection: 200mg tid - duration based on neutropenia/immunosuppression recovery</p> <p>Tx invasive aspergillosis: 200mg po qid then 400mg bid ^{if stable} {if no food 200mg qid}</p>	<p>410 /14 d 3,659 /4wks 3,015-4,519/ 400mg BIDx14-21days</p>
<p>Echinocandins - IV:</p> <p>Caspofungin acetate C</p> <p>Cancidas 50, 70mg vial</p> <p>Micafungin sodium M</p> <p>Mycamine 50mg vial</p> <p>Anidulafungin A</p> <p>Eraxis 100mg vial</p> <p>Broad spectrum; often requiring Infectious Disease Service consult!</p>	<p>C Common: well tolerated;</p> <p>C: fever, phlebitis ^{infusion site}, ↑ALT & AST, histamine-like effects: rash, pruritus, facial swelling</p> <p>M: nausea, vomiting, ↑ALT, AST & ALP</p> <p>A: diarrhea & hypokalemia, ↑ALT</p> <p>Serious:</p> <p>C: hepatotoxicity</p> <p>M: anaphylaxis ^{rare}, febrile neutropenia, hepatic abnormalities, renal insufficiency, hemolytic anemia</p> <p>A: anaphylaxis, hepatic abnormalities, DVT, low BP & flushing (minimize with infusion rate<1.1mg/min)</p>	<p>√ Active: most <i>Candida</i> spp (incl. azole-resistant), <i>Aspergillus</i> spp;</p> <p>C: invasive & esophageal candidiasis; invasive Aspergillosis ^{refractory/intolerant}</p> <p>M: esophageal candidiasis & prevent stem cell transplant invasive candidiasis;</p> <p>A: esophageal candidiasis & candidemia</p> <p>D: ↓ levels of caspofungin: enzyme inducers ie. carbamazepine & rifampin; dexamethasone, efavirenz, nevirapine, phenytoin → consider ↑dose 70mg OD</p> <p>↑caspofungin levels: cyclosporine ↑ hepatic enzymes</p> <p>M: ↑ level of: itraconazole, nifedipine, sirolimus</p> <p>Do not adjust in renal failure; C requires adjustment in liver failure.</p> <p>M: A: LFT's; C: K⁺, LFT's;</p> <p>M: Lytes (K⁺, Mg⁺⁺), Scr, BUN, LFT's, CBC</p> <p>Comment: Preferred for <i>C. Glabrata</i> candidemia ^{IDSA guidelines}</p>	<p>C: Candidemia neutropenic & non-neutropenic: Load: 70mg iv x 1 → 50mg iv once daily Esophageal candidiasis: 50mg iv once daily</p> <p> Liver impairment (Child-Pugh score 7-9): 70mg load → 35mg iv once daily</p> <p>M: Candidemia neutropenic & non-neutropenic: 100mg iv daily; Esophageal candidiasis: 150mg iv daily; Prophylaxis stem cell transplant : 50mg iv daily</p> <p>A: Candidemia neutropenic & non-neutropenic: Load: 200mg iv x1 → 100mg iv od x 14day ^{minimum};</p> <p>Esophageal candidiasis: Load 100mg iv x 1 → 50mg iv od x 14day ^{minimum}</p>	<p>446 /70mg vial 271 /50mg vial 98 /50mg vial 214 /100mg vial</p>
<p>Amphotericin B - IV</p> <p>Amphotericin B deoxycholate (AmBd): Fungizone 50mg vial</p> <p>Lipid formulations:</p> <p>i) Amphotericin B lipid complex (ABLC): Abelcet 100mg vial</p> <p>ii) Liposomal Amphotericin B (L-Am B): Ambisome 50mg vial</p> <p>iii) Amphotericin B colloidal dispersion (ABCD) ^{in US}</p> <p>Infectious Disease consult!</p>	<p>B Common: infusion reactions: fever, chills, shakes, headache, nausea, vomiting, hypotension & tachypnea (worse with early infusions; may pretreat with acetaminophen/NSAID, diphenhydramine & meperidine) ^{70,71}, malaise, weight loss, mild leukopenia, thrombocytopenia</p> <p>Serious: nephrotoxicity (may reduce with Na⁺ loading /lipid formulations), cardiac toxicity, K⁺ & Mg⁺⁺ wasting (may tx with po spironolactone), myopathy ↓liver toxicity ^{lipid formulations}</p> <p>Precautions: nephrotoxic drugs; liposomal amphoB (L-Am-B) has 900mg sucrose/vial -caution diabetes</p>	<p>√ Active against most fungi & protozoa including Zygomycetes; reserved for serious infections; low therapeutic index, ↑↑toxicity; traditional ampho B_{AmBd} preferred tx for severe fungal infections during pregnancy.</p> <p>D: ↑nephrotoxicity: aminoglycosides, cyclosporine, tacrolimus, & other nephrotoxins including chemotherapy</p> <p>↑ toxicity: digitalis ^{low K+}</p> <p>M: CBC, electrolytes K⁺, Mg⁺⁺, liver transaminases ^{if lipid}, renal fx ^{BUN, Scr}</p> <p>Comments: good CNS penetration; lipid formulations: better tolerated, less nephrotoxicity & less infusion reaction problems, but expensive</p>	<p>Dose varies based upon formulation used & indication/organism treated; duration dependent on response; poorly dialyzed.</p> <p>{usual dose range: AmBd: 0.25-1mg/kg/day; Other formulations: 3-5mg/kg/day}</p> <p>☑no longer need for traditional test dose or gradual titration</p> <p>Broad spectrum; often requiring Infectious Disease Service consult!</p>	<p>Fungizone 68 /50mg vial Abelcet 198 /100mg vial Ambisome 121 50mg vial</p>

Extras: **Tinea alba:** sometimes confused with tinea versicolor; non-fungal in origin and does not require treatment beyond usual care for eczema; **Tinea barbae** : fungal infections of the beard area; oral antifungal required.

¹ CPS 2010

² Micromedex 2010

³ Hansten and Horn. Drug interactions

⁴ Medical letter. Treatment guidelines : Antifungal drugs. The Medical Letter 2008(Jan);6(65):1-8. (Medical Letter-Treatment Guidelines-Antifungal drugs. Dec,2009.)

⁵ Mohr J, Johnson M, Cooper T, et al. Current options in antifungal pharmacotherapy. Pharmacotherapy 2008;28(5):614-645

⁶ Pappas PG, Kauffman CA, Andes D, et al: Clinical practice guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. Clin Infect Dis 2009; 48(5):503-535

⁷ Sanford's Guide to Antimicrobial Therapy 2009.

⁸ Crawford F, Hollis S. Topical treatments for fungal infections of the skin skin and nails of the foot. Cochrane Database Syst Rev 2007; (3) CD001434.

⁹ Goldstein AO, Goldstein BG. Onychomycosis. Up-to-date. Accessed 18 August 2009

¹⁰ Olde Hartman TC, van Rijswick E. Fungal Nail Infection. BMJ 2008 Jul 10; 337: a429.

¹¹ Chang CH, Young-Xu Y, Kurth T et al. The Safety of oral antifungal treatments for superficial dermatophytosis and onychomycosis : a meta-analysis. Am J Med 2007 Sep; 120(9):791-8

¹² Crawford F, Young P, Godfrey C, Bell-Syer SE et al. Oral Treatments for toenail onychomycosis: a systematic review. Arch Dermatol 2002 Jun; 138(6):811-6

¹³ Hart R, Bell-Syer SE, Crawford F, Torgerson DJ, Young P, Russell I. Systematic review of topical treatments for fungal infections of the skin and nails of the feet. BMJ. 1999 Jul 10;319(7202):79-82

¹⁴ Gupta AK, Jain HC, Lynde CW, Macdonald P, Cooper EA, et al. Prevalence and epidemiology of onychomycosis in patients visiting physicians' offices: a multicenter canadian survey of 15,000 patients. J Am Acad Dermatol 2000 Aug;43(2 Pt 1):244-8

¹⁵ Sigurgeirsson B; Steingrimsdottir O. Risk factors associated with onychomycosis. J Eur Acad Dermatol Venerol 2004 Jan;18(1):48-51

¹⁶ Pierard GE, Pierard-Franchimont C. The nail under fungal siege in patients with type II diabetes mellitus. Mycoses 2005 Sep;48(5):339-42

¹⁷ Sigurgeirsson B, Olafsson JH, Steinsson JB, et al. Long-term effectiveness of treatment of terbinafine vs itraconazole in onychomycosis : a 5-year blinded prospective follow-up study. Arch Dermatol 2002;138:353-7

¹⁸ Evans EG, Sigurgeirsson B. Double blind, randomized study of continuous terbinafine compared with intermittent itraconazole in treatment of toenail onychomycosis. BMJ 1999;318:1031-35

¹⁹ Warshaw EM, Fett DD, Bloomfield HE, et al. Pulse versus continuous terbinafine for onychomycosis: a randomized, double-blind, controlled trial. J Am Acad Dermatol 2005 Oct;53(4):578-84

²⁰ Natural Medicine Comprehensive Database. Tea tree oil monograph. Accessed 2 October, 2009. (<http://www.naturaldatabase.com/>)

²¹ Pappas PG, Kauffman CA, Andes D, et al. Clinical practice guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. Clin Infect Dis 2009; 48(5):503-535

²² Kauffman CA. Treatment of oropharyngeal and esophageal candidiasis. Up-to-date Accessed 7 August 2009

²³ Gonsalves WC, Stevens Wrightson A, Henry RG. Common oral conditions in older persons. Am Fam Physician. 2008;78(7):845-852

²⁴ Kulak Y, Ankan A, Delibalta N. Comparison of three different treatment methods for generalized denture stomatitis. J Prosthet Dent. 1994 Sep;72(3):283-8

²⁵ Su CW, Jamieson B. Clinical Inquiries. What is the best treatment for oral thrush in healthy infants? J Fam Pract. 2008 Jul;57(7):484-5.

²⁶ Goins RA, Ascher D, Waecker N, et al. Comparison of fluconazole and nystatin oral suspensions for treatment of oral thrush in infants. Pediatric Infect Dis J. 2002;21:1165-67.

²⁷ Leung AK. Gentian violet in the treatment of oral candidiasis. Pediatr Infect Dis J. 1988 Apr;7(4):304-5

²⁸ Walker M Conquering Common Breast-feeding Problems. J Perinat Neonat Nurs 2007;22:267-274

²⁹ Hoddinott P, Tappin D, Wright C. Breast feeding. BMJ 2008;336:881-7

³⁰ Naldi L, Rebora A. Seborrhic dermatitis. N Engl J Med 2009;360:387-96.

³¹ Korting HC, Kiencke P, Nelles S, Rychlik R. Comparable efficacy and safety of various topical formulations of terbinafine in Tinea pedis irrespective of the treatment regimen. Results of a meta-analysis. Am J Clin Dermatol 2007;8(6):357-364

³² Schwärz A. Superficial fungal infections. Lancet 2004;364:1173-82

³³ Goldstein AO, Goldstein BG. Tinea versicolor. Up-to-date. Accessed 21 August 2009

³⁴ Leeming JP, Sansom JE, Burton JL. Susceptibility of Malassezia furfur subgroups to terbinafine. Br J Dermatol 1997 Nov;137(5):764-7

³⁵ Janniger CK, Schwartz RA, Szepletowski JC, et al. Intertrigo and common secondary skin infections. Am Fam Physician 2005; 72:833-8,840

³⁶ Gupta AK, Chow M, Daniel CR, Aly R. Treatments of tinea pedis. Dermatol Clin. 2003 Jul;21(3):431-62.

³⁷ Gupta AK, Chaudhry M, Elewski B. Tinea corporis, tinea cruris, tinea nigra, and piedra. Dermatol Clin. 2003 Jul;21(3):395-400

³⁸ Andrews MD, Burns, M. Common Tinea infections in children. Am Fam Physician. 2008;77(10):1415-1420

³⁹ Sobel JD. Vulvovaginal candidosis. Lancet 2007; 369:1961-1971

⁴⁰ Watson MC, Grimshaw JM, Bond CM, Mollison J, Ludbrook A. Oral Therapy for Recurrent Vulvovaginal Candidiasis. N Engl J Med. 2004 Aug 26;351(9):876-83.

⁴¹ Carter TC, Druschel CM, Romitti PA, et al. Antifungal drugs and the risk of selected birth defects. Am J Obstet Gynecol 2008; 198:191.e1-191.e7

⁴² Sobel JD, Wiesenfeld HC, et al. Maintenance Fluconazole Therapy for Recurrent Vulvovaginal Candidiasis. N Engl J Med. 2004 Aug 26;351(9):876-83

⁴³ Canadian STI guidelines 2008 <http://www.phac-aspc.gc.ca/std-mts/sti-its/guide-lignesdir-eng.php>

⁴⁴ Stary A, Soeltz-Szoets J, Ziegler C et al. Comparison of the efficacy and safety of oral fluconazole and topical clotrimazole in patients with candida balanitis. Genitourin Med. 1996 Apr;72(2):98-102.

⁴⁵ Edwards, SK. European guideline for the management of balanoposthitis. Int J STD AIDS 2001; 12 Suppl 3:68.

⁴⁶ Carter TC, Druschel CM, Romitti PA, et al. Antifungal drugs and the risk of selected birth defects. Am J Obstet Gynecol 2008; 198:191.e1-191.e7

⁴⁷ Sobel JD, Chaim W, Nagappan V, Leaman D. Treatment of vaginitis caused by Candida Glabrata: use of topical boric acid and flucytosine. Am J Obstet Gynecol 2003; 189:1297-1300

⁴⁸ Pirotta M, et al. Effect of lactobacillus in preventing post-antibiotic vulvovaginal candidiasis: a randomised controlled trial. BMJ. 2004 Sep 4; 329 (7465): 548.

⁴⁹ Das Neves J, Pinto E, Teixeira B, et al. Local treatment of vulvovaginal candidosis. Drugs 2008;68(13):1787-1802.

⁵⁰ Scheinfeld N. Diaper dermatitis. A Review and Brief Survey of Eruptions of the Diaper Area. Am J Clin Dermatol 2005; 6(5):273-81

⁵¹ Briggs GG, Freeman RK, Sumner JY. Drugs in Pregnancy and Lactation 7th Edition. Williams & Wilkins, Baltimore, 2008.

⁵² Brüggemann RJ, Alfenaar JW, Blijlevens NM, et al. Clinical relevance of the pharmacokinetic interactions of azole antifungal drugs with other coadministered agents. Clin Infect Dis. 2009 May 15;48(10):1441-58.

⁵³ Chambers WM, Millar A, Jain S, Burroughs AK. Terbinafine-induced hepatic dysfunction. Eur J Gastroenterol Hepatol. 2001 Sep;13(9):1115-8

⁵⁴ Joseph Mo, Pope E. Dermatology, Dipchand A, Friedman J, Bismilla Z, Gupta S, Lam C. The hospital for sick children: Handbook of pediatrics. Elsevier Canada 2009 p.194-207

⁵⁵ Lau, E. Pediatric drug dosing guidelines. In: Dipchand A, Friedman J, Bismilla Z, Gupta S, Lam C. The hospital for sick children: Handbook of pediatrics. Elsevier Canada 2009. p883-996

⁵⁶ Brown SJ. Efficacy of fluconazole for the treatment of onychomycosis. Ann Pharmother 2009;43:1684-91

⁵⁷ Garey KW, Pai MP, Suda KJ, et al. Inadequacy of fluconazole dosing in patients with candidemia based on Infectious Diseases Society of America (IDSA) guidelines. Pharmacoevidemol Drug Saf 2007;16:919-27.

⁵⁸ Janssen Pharmaceutical Products LP. Sporanox (itraconazole) oral solution prescribing information. Titusville, NJ; 2004.

⁵⁹ Hickman JG. A double-blind, randomized, placebo-controlled evaluation of short-term treatment with oral itraconazole in patients with tinea versicolor. J Am Acad Dermatol 1996 May;34(5 Pt 1):785-7

⁶⁰ Kose O, Bulent Tastan H, Riza Gur A, et al. Comparison of a single 400 mg dose versus a 7-day 200 mg daily dose of itraconazole in the treatment of tinea versicolor. J Dermatolog Treat 2002 Jun;13(2):77-9

⁶¹ Fernandez-Nava HD, Laya-Cuadra B, Tiano EAV. Comparison of single dose 400mg versus 10-day 200mg daily dose ketoconazole in the treatment of tinea versicolor. Int J Dermatol 1997;36:64-66

⁶² Goodless DR, Ramos-caro Fa, Flowers FP. Ketoconazole in the treatment of pityriasis versicolor: International review of clinical trial. DICP1991;25:395

⁶³ Mohr J, Johnson M, Cooper T, et al. Current Options in Antifungal Therapy. Pharmacotherapy 2008;28(5):614-645

⁶⁴ Herbrecht R, Denning DW, Patterson TF, et al. Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. N Engl J Med 2002;347:408-15

⁶⁵ Smith J, Safdar N, Knasinski V, et al. Voriconazole therapeutic drug monitoring. Antimicrob Agents Chemother. 2006 Apr;50(4):1570-2

⁶⁶ Balian JD, Sukhova N, Harris JW, et al. The hydroxylation of omeprazole correlates with S-mephenytoin metabolism: a population study. Clin Pharmacol Ther 1995; 57:662-9.

⁶⁷ Spriafil (posaconazole). Pharmacist's Letter/Prescriber's Letter 2007;23(7):230714.

⁶⁸ Zonios DI, Bennett JE. Update on azole antifungals. Semin Respir Crit Care Med. 2008 Apr;29(2):198-210

⁶⁹ Krishna G, Moton A, Ma L, Savant I, et al. of oral posaconazole on the pharmacokinetic properties of oral and intravenous midazolam: a phase I, randomized, open-label, crossover study in healthy volunteers. Clin Ther. 2009 Feb;31(2):286-98

⁷⁰ Pappas PG, Kauffman CA, Andes D, et al. Clinical practice guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. Clin Infect Dis 2009; 48(5):503-535

⁷¹ Goodwin SD, Cleary JD, Walawander CA, Taylor JW, Grasela TH Jr. Pretreatment regimens for adverse events related to infusion of amphotericin B. Clin Infect Dis 1995;20:755-61

⁷² Burks LC, Aisner J, Fortner CL, Wiernik PH. Mependine for the treatment of shaking chills and fever. Arch Intern Med 1980;140:483-4

Other useful reading :

Andrews E, Damle BD, Fang A, Foster G, Crownover P, LaBadie R, Glue P. Pharmacokinetics and tolerability of voriconazole and a combination oral contraceptive co-administered in healthy female subjects. Br J Clin Pharmacol. 2008 Apr;65(4):531-9. Epub 2008 Feb 21

De Berker D. Fungal nail disease. N Engl J Med. 2009;360:2108-16

Donders G, Bellen G, Bytbejter G, et al. Individualized decreasing-dose maintenance fluconazole regimen for recurrent vulvovaginal candidiasis (ReCidIF trial). Am J Obstet Gynecol. 2008 Dec;199(6):613.e1-9. Epub 2008 Oct 30.

Elewski BE, Cáceres HW, DeLeon L, et al. Terbinafine hydrochloride oral granules versus oral griseofulvin suspension in children with tinea capitis: results of two randomized, investigator-blinded, multicenter, international, controlled trials. J Am Acad Dermatol. 2008 Jul;59(1):41-54

Ferwerda B, Ferwerda G, Plantinga TS, et al.. Human dectin-1 deficiency and mucocutaneous fungal infections. N Engl J Med. 2009 Oct 29;361(18):1760-7.

Glocker EO, Hennigs A, Nabavi M, et al. A homozygous CARD9 mutation in a family with susceptibility to fungal infections. N Engl J Med. 2009 Oct 29;361(18):1727-35.

Gupta AK, Bluhm R, Summerbell R. Pityriasis versicolor. J Eur Acad Dermatol Venerol. 2002 Jan;16(1):19-33

Hainer BL. Dermatophyte infections. Am Fam Physician. 2003;67:101-8

Laudenbach JM, Epstein JB. Treatment strategies for oropharyngeal candidiasis. Expert Opin. Pharmacother. 2009; 10(9):1413-1421

Nurbhai M, Grimshaw J, Watson M, Bond C, Mollison J, Ludbrook A. Oral versus intra-vaginal imidazole and triazole anti-fungal treatment of uncomplicated vulvovaginal candidiasis (thrush). Cochrane Database Syst Rev. 2007 Oct 17;4:CD002845. No statistically significant differences were observed in clinical cure rates of anti-fungals administered by the oral and intra-vaginal routes for the treatment of uncomplicated vaginal candidiasis.

Samaranayake LP, Keung Leung W & Jin L. Oral mucosal fungal infections. Periodontology 2000 2009; 49:39-59

Sundar S, Chakravarty J, Agarwal D, Rai M, Murray HW. Single-dose liposomal amphotericin B for visceral leishmaniasis in India. N Engl J Med. 2010 Feb 11;362(6):504-12. RA, Gallagher JC. Drug Fever. Pharmacotherapy. 2010 Jan;30(1):57-69.

Topical treatment of superficial fungal infections. Pharmacist's letter/Prescriber's letter. 2009; (8):250806

Watson MC, Grimshaw JM, Bond CM, Mollison J, Ludbrook A. Oral versus intra-vaginal imidazole and triazole anti-fungal agents for the treatment of uncomplicated vulvovaginal candidiasis (thrush): a systematic review. BJOG. 2002 Jan;109(1):85-95.