

Herpes Zoster Vaccine (ZOSTAVAX)

Bottom Line...

- **ZOSTAVAX** is indicated for the **prevention of shingles in immunocompetent patients age ≥60** May be used for ≥50yr (FDA & NACI). Vaccine efficacy is only about 50-60%.
- **ZOSTAVAX** reduces the risk of **shingles** by 50% (ARR=1.7%, NNT=59) & post-herpetic neuralgia (PHN) by 67% (ARR=0.28%, NNT=364) over 3.1 yrs.

NNT: Eg. for every 364 patients vaccinated with **ZOSTAVAX**, 1 PHN case was prevented & 6 shingles cases were prevented over ~ 3 yrs.

- Efficacy for prevention of shingles is highest in patients **60-69 years old** & decreases with increasing age.
- **ZOSTAVAX** is **not indicated for treating** shingles or PHN, or for preventing primary varicella infection.
- **ZOSTAVAX** use in patients with a **history of shingles** has not been studied. The vaccine can be given, although the precise risk for and severity of shingles is unknown. (A recent episode of shingles may have boosted immunity).
- Cost effectiveness remains to be established. **Cost** per single dose = \$ 175 – 195 given subcutaneously. {Soon **ZOSTAVAX II** stored in fridge & more \$}
- **The risk of shingles ↑ with age, as does the risk for PHN, acute pain & severe rash, however the efficacy of the vaccine declines significantly for PHN after 3 years & 6 years for shingles – so when is the optimal time to vaccinate???** It may be in those 60 to 69 years old.
- **Outstanding Questions: Is ZOSTAVAX safe & effective in immunocompromised patients? Is it beneficial for patients with a history of shingles? What is the long-term effectiveness (will a booster be required)?**

What is ZOSTAVAX? ^{1,2,3,4,5,6,7,8}

- **Herpes Zoster (shingles) vaccine** contains live, attenuated varicella-zoster virus (VZV) (Oka/Merck strain). It is 14 times more potent than **VARIVAX** chickenpox vaccine to induce an immune response to VZV in older adults. It is **not interchangeable** with **VARIVAX**.
- **Shingles** is a common problem (Lifetime incidence=10-30%; up to 50% in those surviving to age 85 & in immunocompromised; not reported to public health; ~ 1 million cases/ year in the USA)
 - It is due to a reactivation of the VZV within the sensory ganglia because of waning cell-mediated immunity. (Rare before age 50.)
 - Symptoms: painful, unilateral vesicular eruption, which usually occurs in restricted dermatomal distribution, rarely crosses the midline.
 - Rash red papules → grouped vesicles → more pustular often around the trunk (lasts 2-3 weeks) gradually crusts over within 7-10day → not infectious; pain precedes the rash in many cases
 - ~ 20% of patients with shingles develop postherpetic neuralgia (PHN) often defined as pain persisting >3 months from the initial onset of the rash; varying severity
 - **Higher risk:** immunosuppressed pts (HIV, Lupus), female, severe rash & pain; **Lower risk:** if African American, infected with wild type virus.
 - Risk of recurrence is 4-7% after 8 years.
 - **↑risk of PHN with ↑age:** incidence of PHN in 3.1 year study: age <60yrs = <2%, 60-69yrs = 7%, >70yrs 19%^{9,10}, **↑acute pain, ↑rash severity.**
 - Shingles & PHN are rarely fatal, but PHN pain can be debilitating, persistent & diminish quality of life. (Differential DX: Herpes simplex, coxsackie, pyoderma)
- Indicated for prevention of shingles in **patients ≥60yrs** ^{FDA ≥50yr}. **Not** for treating shingles, PHN or preventing primary varicella infection.

Is ZOSTAVAX effective? Two Studies: Shingles Prevention Study (SPS)²⁰⁰⁸ & Zostavax Efficacy & Safety Trial (ZEST)²⁰¹²

Shingles Prevention Study (SPS) ^{9,10} : DB RCT, n = 38,546, immunocompetent pts, median age ~69 yr 59-99yr, 59% ♂, 3.1 yr follow-up, excluded those with history of shingles					
Clinical Outcomes at 3.1 years	Vaccine n = 19,270	Placebo n = 19,276	RRR	ARR	NNT/NNH Over 3.1yrs
Incidence of shingles	1.6% n=315	3.3% n=642	51%	1.7%	NNT = 59 (95% CI: 50-72)
Incidence of PHN pain*	0.14 % n=27	0.42 % n=80	67%	0.27%	NNT = 364 (95% CI: 263-589)
≥1 serious adverse event	1.3 % n=255	1.3 % n=254	NS	NS	-
≥1 serious adverse event AE substudy 11	1.9 % n=64 (3345)	1.3% n=41 (3271)	↑53%	↑0.66%	NNH = 152 (95% CI: 79-1692)
Zostavax Efficacy & Safety Trial (ZEST) ^{12,13} : DB RCT, n = 22,439, immunocompetent pts, mean age ~55 yr 50-59 yr, 62% ♀, 1.3 yr follow-up					
Clinical Outcomes at 1.3 years	Vaccine n=11,211	Placebo n=11,228	RRR	ARR	NNT/NNH Over 1.3 yrs
Incidence of shingles	0.27% n=30	0.88 % n=99	↓69.8 %	↓0.61 %	NNT = 164 (95% CI: 142 – 212) <i>Conversion to 3 years NNT = 71</i>
≥1 serious adverse event	0.6% n=69	0.5% n=61	NS	-	-

For every 364 patients vaccinated, 6 cases of shingles & 1 case of PHN is prevented over 3 years.


* pain ≥3 on a scale of 0-10 (0 = no pain & 10 = pain as bad as you can imagine) persisting or appearing ≥90 days after rash onset

- **SPS Short-Term Persistence Substudy¹⁴:** n=14,270 subjects: 7320 vaccine & 6950 placebo followed for an additional 5 years ~8yr follow-up. Efficacy ↓ yearly, **losing statistical significance in the 3rd year post-vaccination for PHN & the 6th year post-vaccination for shingles.** Data from the **SPS Long-Term Persistence Substudy**, which followed subjects for 10 years, is not yet available.
- **Number Needed to Vaccinate (NNV)** is an estimate of the **lifetime risk** of shingles after vaccination. Using Canadian population-based data, assuming vaccination at 65 years of age, the **NNV for ZOSTAVAX is estimated at 11 to prevent one case of shingles & 43 to prevent one case of PHN over the remaining life span of vaccine recipients¹⁵.** Remember: NNT is for a specified time range e.g. vaccinate 59 people to prevent one shingles case & 364 people to prevent one PHN case over 3.1 yrs. ^{SPS} Note: NNV may vastly overestimate the benefit because it assumes that immunity does not wane, an assumption that conflicts with the **SPS** substudy. The most representative values for overall effectiveness are likely somewhere between the 3 year NNT & the lifetime NNV.

What are potential adverse events and drug interactions with ZOSTAVAX? ^{1-3,5,16,17,18,19}

- **Common adverse events include** (compared to placebo):
 - Injection site reactions erythema, pain/tenderness, swelling, pruritis & headache. Most reactions were considered mild in intensity.
 - Post-market reports difficult to establish causal relationship; hypersensitivity incl. anaphylactic reactions; rash; pyrexia; lymphadenopathy injection-site
- **Interactions: Can be administered with other live vaccines give on same day or separate by at least 4 weeks & inactivated vaccines**
 - Must **not** be mixed with any other products in the same syringe. Must be given as separate injections and at different body sites.
 - Can **ZOSTAVAX** be given together with **PNEUMOVAX 23** (pneumococcal vaccine)? Manufacturer says “No”, CDC & PHAC says “Yes”. (An observational study suggests there is no problem with immune response when giving both together. ^{Tseng}) Likely give in separate injection sites.
 - **Co-administration of HZV & pneumococcal vaccine:** currently contraindicated by **ZOSTAVAX** manufacturer **due to concerns about ↓ immunogenicity of HZV** but a large observational study reported no difference in efficacy or safety when **ZOSTAVAX & PNEUMOVAX 23** were administered simultaneously. **Centers for Disease Control (CDC) & Prevention recommends concurrent administration of HZV & pneumococcal vaccines in patients who are eligible for both vaccinations.**


What are other potential cautions regarding the use of ZOSTAVAX? ^{1-3,5}

- **ZOSTAVAX is contraindicated if:** *Consider deferring in acute illness/fever!*
 - Patients have had an anaphylactic or anaphylactoid reaction to gelatin or neomycin contact dermatitis to neomycin is not a contraindication
 - Active untreated tuberculosis or immunocompromised leukemia, lymphoma, neoplasms of the bone marrow/lymphatic system, AIDS/HIV.
 - Immunosuppressive therapy. **Vaccinate ≥14 days prior** (or at least 2 weeks) **starting treatment with immunosuppressives** anti-TNF agents, corticosteroids, etc. Delay administration for at least 1 month after high dose corticosteroids for ≥2 weeks (≥20 mg/day prednisone) or use of anti-TNF agents. ^{ACIP⁰⁸}
- **Can ZOSTAVAX be used in immunocompromised patients?** There is short-term evidence that HZV is safe & effective in patients with certain autoimmune diseases rheumatoid arthritis, psoriasis & inflammatory bowel disease, but there is no data on use patients who may be severely immunocompromised in e.g., HIV/AIDS, leukemia, lymphomas currently undergoing chemotherapy, etc.
 - A recently published retrospective study suggests that HZV in patients taking anti-TNF agents antagonists & other immunosuppressants are effective in ↓ the incidence of shingles & do not pose any additional safety risks²⁰.
 - If post-chemo or post-immunosuppressant, no longer immunocompromised & WBC count ok, wait at least 3 months before giving Zostavax.
 - Acyclovir/famciclovir/valacyclovir should be stopped ≥24 h **before vaccination** & should not be started until 14 days afterward.
- Transmission of virus from vaccine to contacts (e.g. immunosuppressed) not reported, but a theoretical concern.
- Not recommend for patients who received **VARIVAX**. (Patients with hx of zoster can be vaccinated, but may consider **5+ yr delay** to ↑ immune boost effect.)
- Use in age <50 yrs or women of childbearing potential is not recommended. HZV is contraindicated in pregnancy (varicella infection a known fetal risk; no studies). Pregnancy should be avoided for at least 1 month following vaccination. Breast feeding is not a contraindication. 

Is administration of ZOSTAVAX cost effective? ^{2,3,5}

- **ZOSTAVAX** costs \$175 - 195 for single dose. Given the many uncertainties, conclusions about cost-effectiveness remain to be definitively demonstrated. Estimate cost per quality adjusted life-year (QALY) gained \$27,000 - \$112,000 intermediate to high end of acceptable range.

What are the Current Vaccination Recommendations for Herpes Zoster Vaccine (ZOSTAVAX)? ^{21,22,23,24} NACI & ACIP = national advisory committees

-  **NACI** ²⁰¹⁰: recommended for persons ≥60 years of age (Grade A) & may be used for persons 50 – 59 years of age (Grade B)
- **USA** – ACIP ²⁰⁰⁸: routine vaccination for all persons ≥60 yrs; No recommendation for persons <60 yrs
- **History of chicken pox:** HZV can be administered (NACI, Grade A)
- **History of HZ:** patients can be vaccinated. In theory, prior episodes of HZ ↑ immunity & ↓ likelihood of recurrences, but observational evidence is contradictory^{25,26}. A recent study reports the risk of recurrence is ↓ for 12 to 18 months after having HZ so vaccination could be delayed by ≥1 year to take advantage of this natural immunity²⁰.

How is ZOSTAVAX supplied? What is the dosage and how is it administered? ^{1-3,5}

- Supplied in a single-dose vial. Diluent ^{0.7ml} supplied separately. After reconstitution: is a semi-hazy to translucent, off-white to pale yellow liquid (0.65 mL) & contains VZV ≥19,400 PFU (plaque-forming units)
- Prior to reconstitution the vaccine **should be stored frozen** at an average temp of ≤ -15°C may be good for 72hr at up to 8°C, until reconstituted. The diluent should be stored at room temp (20-25°C) or refrigerated (2-8°C). **Administer vaccine immediately after reconstitution, to minimize loss of potency.** Discard if reconstituted vaccine is not used **within 30 mins**. Contains no preservatives (thimerosal free). **{Soon available ZOSTAVAX II stored in fridge & more \$}**
- Individuals should receive a single dose of the entire vial contents, **subcutaneously** deltoid region.
- On the horizon: non-freezer version (**ZOSTAVAX II** – may be ready to market in 2014)²⁷ & non-live vaccine formulations are in development.

Uncertainties

- Of those in the vaccinated group who do get shingles, are severity and complications reduced? Is efficacy retained over longer term?
- As more severe PHN is likely the most important issue, to what extent were the more severe/persistent PHN cases prevented?
 - **Duration of effect & boosters:** Persistence of **ZOSTAVAX** effect beyond 5 years is being studied. The results of this study should help determine the need for revaccination. **No booster dose is recommended at present.**
 - **Does ZOSTAVAX prevent recurrences of shingles after an initial episode?** There are no anticipated safety concerns but no studies have investigated efficacy.
 - **Will people who have received varicella vaccine be at risk of shingles as they age?** Currently, it is thought that varicella vaccination ↓ the risk of severe shingles but it is not known whether this effect will persist as people age.

Shingles Extras ^{27,28}:

- Antivirals (e.g. valacyclovir 1g TID or acyclovir 800mg 5x/day) x7 days \$70; effective in shingles treatment for age >50 if used within 24-72hrs of rash onset.
- See RxFiles Chronic Non-Cancer Pain chart for PHN pain treatment (9th Ed, pg 67) → e.g. nortriptyline, gabapentin, opioid, capsaicin.
- See RxFiles Adult Vaccines Chart (9th ed, pg 50).

Additional articles:

- Breuer J, Pacou M, Gauthier A, et al. Herpes zoster as a risk factor for stroke and TIA: A retrospective cohort study in the UK. *Neurology*. 2014 Jan 21;82(3):206-12
- Cohen JI. Clinical practice: Herpes zoster. *N Engl J Med*. 2013 Jul 18;369(3):255-63.
- Irwin MR, Levin MJ, Laudenslager ML, et al. Varicella Zoster Virus-Specific Immune Responses to a Herpes Zoster Vaccine in Elderly Recipients With Major Depression and the Impact of Antidepressant Medications. *Clin Infect Dis*. 2013 Feb 13.
- Keating GM. Shingles (herpes zoster) vaccine (zostavax®): a review of its use in the prevention of herpes zoster and postherpetic neuralgia in adults aged ≥50 years. *Drugs*. 2013 Jul;73(11):1227-44.
- Kolber MR, Korownyk C, Nickonchuk T. Zoster vaccine. *Can Fam Physician*. 2013 Feb;59(2):157.
- Langan SM, Smeeth L, Margolis DJ, Thomas SL. Herpes zoster vaccine effectiveness against incident herpes zoster and post-herpetic neuralgia in an older US population: a cohort study. *PLoS Med* 2013;10:e1001420.
- Langan SM, Minassian C, Smeeth L, Thomas SL. Risk of Stroke Following Herpes Zoster: A Self-Controlled Case-Series Study. *Clin Infect Dis*. 2014 Apr 2.
- Schmader KE, Oxman MN, Levin MJ, et al. Shingles Prevention Study Group. Persistence of the efficacy of zoster vaccine in the shingles prevention study and the short-term persistence substudy. *Clin Infect Dis*. 2012 Nov 15;55(10):1320-8.
- Tseng HF, Smith N, Sy LS, Jacobsen SJ. Evaluation of the incidence of herpes zoster after concomitant administration of zoster vaccine and polysaccharide pneumococcal vaccine. *Vaccine*. 2011 May 9;29(20):3628-32
- Yawn BP, Gilden D. The global epidemiology of herpes zoster. *Neurology*. 2013 Sep 3;81(10):928-30.

References: Herpes Zoster Vaccine (ZOSTAVAX)

- Merck Frosst Canada Ltd. Product Monograph: Zostavax™. Available @: http://www.merckfrosst.ca/assets/en/pdf/products/ZOSTAVAX-09_08-a_129172-E.pdf (accessed Sept 2009).
- Herpes zoster (Zostavax) vaccine. Pharmacist's Letter/Prescriber's Letter 2006;22(7):220702.
- Singh A, England K. Q: Who should receive the shingles vaccine? *Cleve Clin J Med*. 2009 Jan;76(1):45-8.
- Albrecht MA. Clinical manifestations of VZV infection: Herpes zoster. In: UpToDate Online. UpToDate Waltham, MA. Available from: www.uptodate.com (accessed September 2009).
- Harpaz R, et al. Prevention of herpes zoster: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2008 Jun 6;57(RR-5):1-30.
- Liesegang TJ. Varicella zoster virus vaccines: effective, but concerns linger. *Can J Ophthalmol*. 2009 Aug;44(4):379-84
- Bajwa ZH, Warfield CA, Crovo DG. Postherpetic neuralgia. In: UpToDate Online. UpToDate Waltham, MA. Available from: www.uptodate.com (accessed September 2009).
- Jung, Beth F., Johnson, Robert W., Griffin, David R.J., Dworkin, Robert H. Risk factors for postherpetic neuralgia in patients with herpes zoster *Neurology* 2004 62: 1545-1551.
- Oxman MN et al for the Shingles Prevention Study Group. A vaccine to prevent herpes zoster and postherpetic neuralgia in older adults. *N Engl J Med*. 2005 Jun 2;352(22):2271-84.
- Fekete T. A varicella-zoster virus vaccine reduced the burden of illness of herpes zoster in older adults. *ACP J Club*. 2005 Nov-Dec;143(3):61.
- Simberkoff MS, Arbeit RD, Johnson GR, et al. Shingles Prevention Study Group. Safety of herpes zoster vaccine in the shingles prevention study: a randomized trial. *Ann Intern Med*. 2010 May 4;152(9):545-54.
- Schmader K, Levin M, Gnann J et al. Efficacy, safety, and tolerability of herpes zoster vaccine in persons aged 50–59 years. *Clin Infect Dis* 2012;54:922–8.
- Li H, Lu S, Liu J. Herpes zoster vaccination in people aged 50–59 years. *Clin Infect Dis* 2012;54:929-30.
- Schmader K, Oxman M, Levin M et al. Persistence of the efficacy of zoster vaccine in the shingles prevention study and the short-term persistence substudy. *Clin Infect Dis* 2012.
- Brisson M. Estimating the number needed to vaccinate to prevent herpes zoster-related disease, health care resource use and mortality. *Can J Public Health*. 2008 Sep-Oct;99(5):383-6.
- Tseng HF, Smith N, Sy LS, Jacobsen SJ. Evaluation of the incidence of herpes zoster after concomitant administration of zoster vaccine and polysaccharide pneumococcal vaccine. *Vaccine* 2011;29:3628-32.
- Shapiro M, Kvern B, Watson P et al. Update on herpes zoster vaccination. *CFP* 2011;57:1127-31.
- CDC. Herpes zoster vaccination for health care professional. June 28, 2012. Available at <http://www.cdc.gov/vaccines/vpd-vac/shingles/hcp-vaccination.htm>. Accessed October 2012.
- National Advisory Committee on Immunization (NACI). Canadian immunization guide. 7th ed. Ottawa, ON: Public Health Agency of Canada; 2006. Available from: <http://origin.phac-aspc.gc.ca/publicat/cig-gci/index-eng.php>. Accessed September 2012.
- Zhang J, Fenglong X, Delzell E. Association between vaccination for Herpes zoster and risk of Herpes zoster infection among older patients with selected immune-mediated diseases. *JAMA*. 2012;308(1):43-49.
- National Advisory Committee on Immunization (NACI). Advisory. Statement on the recommended use of herpes zoster vaccine. *Can Commun Dis Rep* 2010;36(ACS-1):1-19. Available at <http://www.phac-aspc.gc.ca/publicat/ccdr-mtc/10pdf/36-acs-1.pdf>. Accessed September 2012.
- ACIP 2008 Prevention of Herpes Zoster Recommendations of the Advisory Committee on Immunization Practices (ACIP) June 6, 2008 / 57(05):1-30 http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5705a1.htm?cid=rr5705a1_e
- Harpaz R, et al. Prevention of herpes zoster: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2008 Jun 6;57(RR-5):1-30.
- Update on Herpes Zoster Vaccine: Licensure for Persons Aged 50 Through 59 Years. *Morbidity and Mortality Weekly Report (MMWR)* November 11, 2011 / 60(44):1528-1528 <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6044a5.htm>.
- Yawn BP, Wollan PC, Kurland MJ, St Sauver JL, Saddier P. Herpes zoster recurrences more frequent than previously reported. *Mayo Clin Proc*. 2011 Feb;86(2):88-93. (Shingles ~6% at 8yrs)
- Tseng HF, Chi M, Smith N, et al. Herpes Zoster Vaccine and the Incidence of Recurrent Herpes Zoster in an Immunocompetent Elderly Population. *J Infect Dis*. 2012 Jun 4.
- PL Detail-Document 280911, Treatment of Shingles. Pharmacist's Letter/Prescriber's Letter. January 2012.
- Fashner J, Bell AL. Herpes Zoster and Postherpetic Neuralgia: Prevention and Management. *Am Fam Physician*. 2011 Jun 15;83(12):1432-1437.

We would like to acknowledge the following contributors and reviewers: Dr. S. Sanche (SHR-Infectious Disease), Y Shevchuk (UofS College of Pharmacy), D. Dr. B. Tan (member NACI; SHR-Ped) & the RxFiles Advisory Committee. Prepared by Jason Keilly (PharmD-UofTy), Loren Regier BSP, BA, Brent Jensen BSP, Karen Jensen MSc BSP, Debbie Bunka BScPharm, Lynette Kosar MSc BSP, Julia Bareham BSP BSc

DISCLAIMER: The content of this newsletter represents the research, experience and opinions of the authors and not those of the Board or Administration of Saskatoon Health Region (SHR). Neither the authors nor Saskatoon Health Region nor any other party who has been involved in the preparation or publication of this work warrants or represents that the information contained herein is accurate or complete, and they are not responsible for any errors or omissions or for the result obtained from the use of such information. Any use of the newsletter will imply acknowledgment of this disclaimer and release any responsibility of SHR, its employees, servants or agents. Readers are encouraged to confirm the information contained herein with other sources.

Copyright 2012 – RxFiles, Saskatoon Health Region (SHR) www.RxFiles.ca