Nov 2025

Bottom line: These anti-amyloid medications generally achieved statistical significance based on their primary outcomes, however, they did not appear to reach a proposed clinically meaningful benefit. The harms (including ARIA-H/E) and the need for regular MRI monitoring limit practical application and add to overall cost of treatment. These novel drugs pose a theoretical disease-progression-altering capacity based on amyloid plaque reduction; however, their clinical value and long-term outcomes remain unclear.

76-week tx duration	TRAILBLAZER-ALZ 2	CLARITY AD	EMERGE and ENGAGE Identically Designed RCTs
for each	Donanemab KISUNLA (2023)	Lecanemab LEQEMBI (2023)	Aducanumab ADUHELM (2022)
Population Highlights	 Included patients with Alzheimer's disease (AD), with mild cognitive impairment or with mild dementia Mean age across RCTs: ~70-73yr MMSE across RCTs needed to be in the upper range: ~20-30 (higher score means less impairment; max score on scale = 30) Population was selective: ~60-75% screened out due to inclusion/exclusion criteria 		
Intervention / Comparator	Donanemab 1400mg (with 700mg dose titration x first 3 doses) vs placebo IV q4 weeks for 76 weeks	Lecanemab 10mg/kg vs placebo IV q2 weeks for 76 weeks	Aducanumab low dose (3 or 6mg/kg target dose) vs aducanumab high dose (10mg/kg target dose) vs placebo IV q4 weeks for 76 weeks
Outcomes	 See individual RxFiles Trial Summaries for more details. RCTs achieved statistically significant results in 1° outcome however, differences seen did not meet what would usually be considered the minimal clinically important difference (MCID) 1 RCT (EMERGE) had statistically significant results 1 RCT (ENGAGE) had non-significant results Neither RCT achieved a clinically significant change in 1° outcome 		
CDR-SB (An 18-point scale to assess cognition & function; lower score = ↓ impairment)	CDR-SB (2º outcome): - 0.68 point difference vs placebo	CDR-SB (1º outcome) - 0.45 point difference vs placebo	CDR-SB (1º outcome) – high dose EMERGE: - 0.39 point difference vs placebo ENGAGE: + 0.03 point difference vs placebo
	The commonly suggested minimally important difference (MCID) on the CDR-SB scale is 1 point. None of the RCTs achieved this.		
iADRS (A 144-point scale to assess cognition & function; higher score =	 iADRS (1° outcome): + 2.92 point difference vs placebo to assess cognition & function (calculated as composite of ADCS-iADL + ADAS-Cog 13 scoredid not achieve pre-determined minimum important difference (MCID in AD with mild cognitive impairment=MCID in AD with mild dementia=9) 	not applicable not reported as a composite.	
Other	Other cognitive assessments conducted showed similarly small differences in score.	Similar relative improvements seen in other measures of cognition and functioning.	Conflicting results between 2 RCTs in measures of cognitive and functional improvement; small, statistically significant improvement (not above the threshold for clinically significant benefit) for high dose in EMERGE .
Amyloid Plaque Reduction	All RCTs reported a reduction in amyloid plaques, which is of interest given the proposed pathophysiology of Alzheimer's disease. However, this is a surrogate endpoint of interest compared to the clinical endpoints above of relatively more importance.		
3.00.000	All RCTs had adverse events (AE) of concern.		
Notable Harms	ARIA-H and ARIA-E (ARIA=amyloid related imaging abnormalities, ARIA-H=ARIA with microhemorrhages, ARIA-E=ARIA with edema): NNH≈3 to 12/76wk. Other AE of interest included infusion reactions, falls, & reduction in brain volume. Serious Adverse Events (SAE) were more common in treated group than placebo (NNH≈37 to 62/76wk), as were discontinuation rates.		
Additional Considerations	Diagnostic supports (PET Scans and MRI) wer monitoring for harm When analyzing the Kaplan Meier Curves fo the placebo group by abo	Drug Cost: \$26,500 USD/year - FDA approved 2023 - HC approved (with conditions) Oct 2025, not yet marketed as of Nov 2025 - Requires imaging which adds to the burden & costs imposed on the health care system (relative to the apparent lack of clinically important le part of the protocols and required for therapy as pathroughout treatment, adding to the burden on finar the RCTs, one way to look at the potential benefit is ut 2 months. This, with the caveat that the difference	art of diagnostic criteria for trial enrollment as well as incial and human resources. It that the decline in scores for the treated group lags agained is still less than the MCID. It is be noticeable. Yet, there may be potential for some

L Regier, BSP BA © www.RxFiles.ca

Nov 2025

Acknowledgements: Written by Loren Regier, as a synopsis of the overview and trial summaries created by M Kurz, E Holmlund, and N Helson who were all PharmD Candidates when the contributed this activity in 2023. Finalized by Amy Wiebe in 2025.

Other Reviewers: Erin Yakiwchuk, Tina Korownyk.

Disclosures: No conflicts of interest are reported by the authors.

Disclaimer: RxFiles Academic Detailing is part of the College of Pharmacy and Nutrition at the University of Saskatchewan. The content of this work represents the research, experience and opinions of the authors and not those of the University of Saskatchewan. Neither the authors nor the University of Saskatchewan nor any other party who has been involved in the preparation or publication of this work warrants or represents that the information contained herein is accurate or complete, and they are not responsible for any errors or omissions or for the result obtained from the use of such information. Any use of the materials will imply acknowledgment of this disclaimer and release any responsibility of the University of Saskatchewan, its employees, servants or agents. Readers are encouraged to confirm the information contained herein with other sources.

References for ANTI-AMYLOID MEDICATION FOR ALZHEIMER'S DISEASE: Overview of Landmark Trials

- 1. Budd Haeberlein, S., Aisen, P., Barkhof, F. et al. Two Randomized Phase 3 Studies of Aducanumab in Early Alzheimer's Disease. J Prev Alzheimers Dis 2022;9(2):197–210. https://doi.org/10.14283/jpad.2022.30.
- 2. Sims JR, Zimmer JA, Evans CD, et al. Donanemab in Early Symptomatic Alzheimer Disease: The TRAILBLAZER-ALZ 2 Randomized Clinical Trial. JAMA. 2023;330(6):512–527. doi:10.1001/jama.2023.13239.
- 3. Van Dyck CH, Swanson CJ, Aisen P, et al. Lecanemab in Early Alzheimer's Disease. N. England J.M. 2023; 388(1): 9-21. doi:10.1056/NEJMoa2212948.
- 4. Biogen Canada. Biogen Canada announces intention to transition Aduhelm access program in Canada [Press release]. (2022, June 9). Available from: https://www.biogen.ca/en-ca/news/2022-06-09-news.html.