

**Critical Appraisal of Drug Studies** <sup>6,7</sup>

**A) Is the study valid?**

1. Were patients **randomized** to treatment (tx) groups & was **allocation concealed** (AC)? {Without concealment, 37% bias in favor of tx. Sealed, opaque envelopes or central registry used to attain AC<sup>8,9</sup>}
2. Was everyone (patients, physicians, investigators, assessors) **blinded** to tx? {Especially important for assessors of subjective outcomes e.g. Pain.}
3. Was the study **controlled**? (e.g. RCT: inclusion of placebo or active control group/arm; in an "N of 1" trial, patient is their own control.)
4. Were treatment & control **groups similar** in prognostic factors for outcome of interest at beginning of study? If not, were adjustments made?
5. Were **all patients accounted for** at end? {Missing patients addressed?}
6. Was data analyzed based on groups patients were initially randomized to? {Intention to treat or **ITT**; protects integrity of prognostic randomization; per protocol (PP) analysis may also be of interest (e.g. non-inferiority trials)}
7. Were **groups treated similarly** except for study intervention?
8. How was the study **funded** (role of funder)? Was study **stopped early**?
9. Was study type, design & comparator drug & dose a good choice?

**B) What are the study results?**

1. What was the primary (1°) endpoint? What were the secondary (2°) endpoints? Were endpoints & subgroups pre-specified?<sup>10</sup> Avoid data mining!
2. What was the difference in **outcomes**? (Both benefits & harms.)
3. Were the differences **statistically significant**? ... **clinically significant**? {What were the 95% confidence intervals (CIs) or p values? Does the CI cross line of no effect?}
4. What are the **absolute** and **relative risk reductions**, or increases?
5. What is the number needed to treat (**NNT**) &/or harm (**NNH**)?

**C) Does this study matter to my patients?**

1. How **clinically relevant/important** are the outcomes?
2. Were the patients similar to those in my practice? {Consider **inclusion & exclusion criteria**: very sick, old, young, drug interactions & complicated/co-morbid patients often excluded.}
3. Do treatment **benefits outweigh the risks**, costs & impact on life?

**Types of Studies** (from low to high level of evidence) <sup>11</sup>

- **Case-control study**: a retrospective observational study which selects patients a) with the outcome of interest (cases) & b) without that outcome (controls); attempts to find exposures linked to the outcome.
- **Cohort study**: an observational study in which 2 groups (cohorts) are observed over time for an outcome. One cohort has exposure to a condition/treatment that the other does not. {Observational studies: association does not prove causation!  
Strength of association: RR: 1.01-1.5 *weak*; 1.51-3 *moderate*; >3 *strong*.<sup>12</sup>}
- **Randomized controlled trial (RCT)**: a prospective study in which patients are randomized to treatment or control groups (equal chance at being assigned to any group, like the flip of a coin). Groups are followed for the outcome of interest.
- **Crossover RCT**: a design in which each patient receives both treatments in two phases separated by a washout period. Each patient serves as own control, thus less variability in outcomes, & smaller sample size OK; period effects may limit findings
- **Systematic Review (SR)**: a systematic collection, review & presentation of available studies addressing a clinical question using specific criteria & methods; may, or may not, include meta-analysis. e.g. *Cochrane*<sup>13</sup>/*Campbell*<sup>14</sup>/*CADTH Reviews*<sup>15</sup>

{**Meta-analysis**: the combining of studies meeting prespecified criteria, addressing a clinical question. Results are calculated from each study's data, then pooled. ↑ sample size & statistical power useful if single trial or subgroup analysis underpowered., **Assess appropriateness** of a) variables & outcomes; b) studies included; c) if study quality & heterogeneity accounted for.}

[**Level of evidence for tx**: SR>RCT>observational study>expert opinion.<sup>15</sup>  
Observational studies useful for safety, generalization to different populations, & insights into real world effect, especially when specific RCT not practical.<sup>38</sup>]

**Caution**: Lots of low quality RCTs may not be better than 1 good quality RCT!  
A low quality SR, or a SR of low quality trials does not constitute high-level evidence.]

**Terms: Related To Validity**

- **Bias**: design flaws leading to over/underestimation of treatment effect e.g. recall bias, selection bias, publication bias; confounding factors esp observational studies
- **Blinding**: if investigators, patient etc are unaware of who receives tx vs control, they are less likely to inappropriately report better results with tx.

{**CONSORT Statement** a checklist of standards for standardized reporting of RCTs intended to reduce bias. <http://www.consort-statement.org/>}

**Study Results: Size Of The Treatment Effect** <sup>16,17,18,19</sup>

- **Event rate (ER)**: the number of people experiencing the event as a proportion of total number of people in the population or group  
-Experimental ER (**EER**): {# events in experimental group / total in exp. group}  
-Control group ER (**CER**): {# events in control group / total in control group}
- **Relative risk (RR)** or risk ratio: {EER/CER}
- **Relative risk reduction (RRR)**: the RR subtracted from 1 {RRR=1-RR} [Whereas ARR varies with type of population treated, RRR is often more constant.]
- **Absolute risk reduction (ARR)**: the arithmetic difference between the 2 event rates {CER-EER} [If ↑ risk: **ARI**= absolute risk increase]
- **Number needed to treat (NNT)**: the number of people who would have to be treated with the studied intervention for the studied time period to see 1 extra of them benefit. {**NNT=100/ARR%**}
- **Number needed to harm (NNH)**: number of people who would have to be treated with the studied intervention for the studied time period for 1 extra person to experience the adverse event. {**NNH=100/ARI%**}
- **Odds ratio (OR)**: = experimental event odds / control event odds; especially used in case-control studies where baseline risk is not known; also used in meta-analysis. When events are rare, the OR is similar to the RR; however, OR rate exaggerated relative to RR when events more common. {Link [www.cebm.net/](http://www.cebm.net/): tool for converting OR to NNT<sup>20</sup>}
- **Point estimate**: the trial result used as best estimate of the true effect
- **Hazard ratio (HR)**: like RR but more accurate; accounts for the time each participant was in the study before having event or withdrawing.

**Study Results: Precision of Treatment Effect** <sup>21</sup>

- **Confidence Interval (CI)**: a 95% CI provides the range of values we are 95% certain that overlap the true value. CI's indicates the precision of the estimate; where CI's are wide, they indicate less precise estimates of effect (just an estimate of the worst & best case scenario of the outcome; related to p-value) {For ratios, a CI that includes 1 means possibility of no difference. For ARR, ARI, NNT, NNH, a CI that includes zero means possibility of no difference between tx. Non-significant results, *trends*, may provide clues re uncertainties & future research.}
- **Type 1 (or α) error**: the *false positive*; to find a difference when there is none. **p-value**: reflects type 1 error. A p <0.05 suggests a <1 in 20 probability that any difference is due to chance (statistically significant by convention). The smaller the **p-value**, the less likely that the result is due to chance.
- **Type 2 (or β) error**: the *false negative*; to conclude there is no difference when there really is a difference (e.g. if not enough patients enrolled)
- **Heterogeneity**: when studies within a meta-analysis have more variation than expected; may indicate inappropriate to combine studies.<sup>22</sup> {**Q statistic**: measure of within-study variance; **I<sup>2</sup>**: ratio of variability among studies to total variation.}

**Calculations Example: 1 yr trial**

- 200 patients in Control group
- 200 patients in treatment (tx) group
- Deaths: Control grp: 40. **CER=40/200=0.2**  
tx grp: 30. **EER=30/200=0.15**

**RRR**  
= (0.20 - 0.15)/0.20 X 100  
= 25% {risk of event is reduced by 25%}

**ARR**  
20% - 15% = 5%  
{absolute risk of event is reduced by 5%}

**NNT**  
= 100/5%  
= 20

**NNH**: if 60% of patients in tx group experienced *headaches* compared with 27% in control group (ARI=33%)  
**NNH=100/33% = 3**

For every 20 patients treated for 1yr. there is 1 less *death*: & for every 3 patients treated there will be 1 extra *headache*.

**A few NNTs / NNHs of interest**

	NNT
↓ mortality with simvastatin 20-40mg/day over 5.4yrs vs placebo in patients with CHD <sup>45</sup>	30 / 5.4yrs
↓ mortality with metformin 2550mg/day over 10 years vs non-intensive tx in obese T2DM patients UKPDS-34	14 / 10 yrs
↓ CV death/MI/stroke; clopidogrel 75mg/day + ASA vs ASA alone in ACS pts (↑ bleeding: <b>NNH=99</b> ) <b>CURE</b>	48 / 9mo
↓ neuropathic pain by ≥50% vs placebo: TCAs ~75mg/day, gabapentinoids, SNRIs duloxetine 60mg/day; (short-term) <sup>37</sup>	4, 7, 8

**What makes for a good NNT?** *It all depends!!!*  
NNTs will vary greatly with variations in baseline population risk, duration of tx, & type & number of endpoints included in composite. The value of the endpoint also varies from patient to patient.

**How do the results matter to me, my patients & society?**

- **Clinical significance vs statistical significance**: some studies may detect extremely small statistically significant differences between groups; however magnitude of effect (e.g. **NNT**) may be too small to change practice. Evaluate both 1) the endpoint, & 2) the **NNT** or **NNH**. {e.g. small cognitive score improvement not noticeable to patient.<sup>23,24</sup>}
- **Composite endpoints**: combining endpoints can increase a study's power allowing for smaller or shorter trials. Outcomes should have **similar value**. Examination of individual outcomes can be important in interpretation as one endpoint may be the primary *driver*. {e.g. In **DREAM**, outcome of "diabetes diagnosis <sup>the driver</sup> or death" = example of unequal endpoints.<sup>25</sup>}
- **Surrogate endpoints**: an endpoint meant to reflect / be correlated with another endpoint (e.g. BP/LDL/A1c for CV events; CD4 cell count for HIV mortality). **Clinical outcomes are more important** since surrogate endpoints **assume** correlation with an outcome which may, or may not always be true.<sup>26</sup> {eg. lower A1c target <sub>≤6%</sub> **ACCORD**: but ↑ death; doxazosin ↓ BP **ALLHAT** but ↑ HF/stroke; & clofibrate **WHO-CLOF** ↓ LDL but ↑ death.}
- **Other considerations**: What uncertainties remain, & how should they be weighed (e.g. legitimate vs illegitimate uncertainty<sup>39</sup>)? Has the drug been studied enough to detect rare serious adverse events (SAE)? What duration is studied & what are the potential benefits/harms over a longer term of exposure? Is real-world experience consistent with clinical trial data? Any insights from subgroup analysis (see **ICEMAN tool**<sup>40</sup>)? Cost? How benefits & harms are described e.g. RR vs **NNT** will also affect decisions.<sup>27</sup>
- **What patient specific &/or societal values need to be considered?**

**Heads Up! Know what the numbers are telling you.**

⇒ You "double" your chance of winning a lottery if you buy a 2<sup>nd</sup> ticket; however your chance of winning is impacted more by whether 2 tickets or 2 million tickets are sold!

**Beware of the Relatives** 😊

- Benefits are often given as **relative** numbers, whereas harms are often given as **absolute** numbers. This tends to exaggerate benefits & minimize the harms. ⇒ **Look for NNTs & NNHs.**  
{e.g. **Vioxx** monograph 2004<sup>41</sup>: reported ~ 50% ↓ in GI complications with Vioxx 50mg/day vs naproxen 500mg BID & a thrombotic event rate of 1.8% (Vioxx) vs 0.6% (naproxen). Actual **GI complication** reductions 0.59% vs 1.37% (ARR=0.78; **NNT=129**); whereas **thrombotic risk** worse (**NNH=83**). **VIGOR** }  
{e.g. Oral contraceptives: risk of DVT in a younger, non-smoking ♀ may be ↑ 300% but absolute risk is <1.5/10,000 /yr, & lower than risk in pregnancy}
- **Non-Equivalent Durations & Risk/Benefit Perception**  
- Benefits are often given for total duration of trial which may be several years, whereas **harms are often given per year**.  
{e.g. **UKPDS-33**: aggressive glucose control benefit on microvascular endpoints given per **10 years**; risks of hypoglycemia were given **per year**.<sup>28</sup>}
- **Analysis: Pooling Together or Dividing Out**  
- Discussing the multiple benefits of a composite endpoint while individually sorting out harms may **minimize risk perception**.  
{e.g. In **WHI**, risk of just breast ca with HRT was **8/10,000 pt-years**; yet risk of

## Other Online EBM Resources/Links:



EBM Portal **Links** (SK): [http://web.mac.com/malees/Primary\\_Care\\_Portal/EBM.html](http://web.mac.com/malees/Primary_Care_Portal/EBM.html); Evidence Updates service: <http://plus.mcmaster.ca/EvidenceUpdates/Default.aspx>

**General:** U of T: <http://www.cebm.utoronto.ca/>; Oxford: <http://www.cebm.net/?o=1011>; McMasters: How to teach evidence based clinical practice – **Links:** <http://hsl.mcmaster.ca/ebcp/>. Dynamed: [www.ebscohost.com/dynamed/](http://www.ebscohost.com/dynamed/)  
User's Guide: UoFA, Centre for Health Evidence: <http://www.cche.net/usersguides/main.asp>; UBC: <http://www.ti.ubc.ca/>; Grey Literature Searching: <http://www.cadth.ca/index.php/en/cadth/products/grey-matters>  
SchHARR Intro to Evidence Based Practice (Sheffield, UK) <http://www.shf.ac.uk/scharr/ir/netting/>; BMJ – Clinical Evidence **Links:** [http://clinicalevidence.bmj.com/ceweb/resources/useful\\_links.jsp](http://clinicalevidence.bmj.com/ceweb/resources/useful_links.jsp); NNTs <http://www.thennt.com/>  
**Clinical significance CALCULATORS:** UBC: <http://spph.ubc.ca/sites/healthcare/files/calculators/significance.html>; Wisconsin: <http://intsmain.is.mcw.edu/clinical/bayes.html>. Essential Evidence Plus: <http://www.essentialevidenceplus.com/>  
Dalhousie **Katie** Clinical Significance Calculator: <http://ktcalc.cme.dal.ca/site/login.php> Z-score: <http://www.socscistatistics.com/pvalues/normaldistribution.aspx> **Teaching EBM Videos:** McMaster **Guyatt:** [http://ebm.mcmaster.ca/materials\\_videos.htm](http://ebm.mcmaster.ca/materials_videos.htm)  
**Other tools of interest:** **ICEMAN** - to assess subgroup heterogeneity effect <sup>40</sup>

## RxFiles – Select Trial Summaries (more available online at [www.RxFiles.ca](http://www.RxFiles.ca))

**Anemia:** Trials Summary: <http://www.rxfiles.ca/rxfiles/uploads/documents/CHT-Anemia-Key-Trials.pdf>  
**Asthma:** Asthma Landmark Trials: Treatment of "Mild" or Intermittent Adult & Adolescent Asthma <sup>2021</sup>; Asthma Trials/SR Overview/Summary <sup>2006</sup>  
Novel-START: <https://www.rxfiles.ca/RxFiles/uploads/documents/members/ts-NovelStart.pdf> <sup>2021</sup>  
PRACTICAL: <https://www.rxfiles.ca/RxFiles/uploads/documents/members/ts-PRACTICAL.pdf> <sup>2021</sup>  
SYGMA-1: <https://www.rxfiles.ca/RxFiles/uploads/documents/members/ts-SYGMA1.pdf> <sup>2021</sup>  
SYGMZ-2: <https://www.rxfiles.ca/RxFiles/uploads/documents/members/ts-SYGMA2.pdf> <sup>2021</sup>  
**Dementia:** CATIE-AD: <http://www.rxfiles.ca/rxfiles/uploads/documents/Psych-CATIE-AD-trial-summary.pdf>  
**Diabetes:** Landmark Trials Summary: Glucose: <http://www.rxfiles.ca/rxfiles/uploads/documents/CHT-Diabetes-Landmark-Trials-Links.pdf>  
Landmark Trials Summary: NON-Glucose: <http://www.rxfiles.ca/rxfiles/uploads/documents/members/CHT-DIABETES-Landmark-Trials-Non-Glucose.pdf>  
ACCORD-ADVANCE Comparison: <http://www.rxfiles.ca/rxfiles/uploads/documents/Diabetes-A1C-ACCORD-vs-ADVANCE-COMPARISON.pdf>  
ACCORD-BP & LIPID: <http://www.rxfiles.ca/rxfiles/uploads/documents/ACCORD-BP-Lipid-Trial-Overview.pdf>  
ACCORD: Glucose <http://www.rxfiles.ca/rxfiles/uploads/documents/Diabetes-Targets-ACCORD-A1C.pdf>  
ADVANCE: <http://www.rxfiles.ca/rxfiles/uploads/documents/Diabetes-ADVANCE-trial.pdf>  
AVANDIA & CV risk – Meta-analysis: <http://www.rxfiles.ca/rxfiles/uploads/documents/Diabetes-Avandia-CV-Meta-Comments.pdf>  
DREAM: <http://www.rxfiles.ca/rxfiles/uploads/documents/Dream-QandA.pdf>  
ELIXA: Lixisenatide : <http://www.rxfiles.ca/rxfiles/uploads/documents/Lixisenatide-ELIXA%20Trial%20Summary.pdf> <sup>2016</sup>  
EMPA-REG: <http://www.rxfiles.ca/rxfiles/uploads/documents/EMPA-REG%20Trial%20Summary.pdf> <sup>2016</sup>  
LEADER: Liraglutide: <http://www.rxfiles.ca/rxfiles/uploads/documents/Leader-Liraglutide%20VICTOZA%20Qand%20Cardiovascular%20Outcomes%20in%20Type%202%20Diabetes.pdf> <sup>2016</sup>  
RECORD: <http://www.rxfiles.ca/rxfiles/uploads/documents/Diabetes-RECORD-Trial-Summary.pdf>  
SAVOR-TIMI 53: <http://www.rxfiles.ca/rxfiles/uploads/documents/SAVOR-TIMI-53-Saxagliptin-CV-Outcomes-Trial-Summary.pdf>  
TECOS: Sitagliptin CV outcomes: <http://www.rxfiles.ca/rxfiles/uploads/documents/TECOS-Trial-Summary.pdf> <sup>2016</sup>  
**Hypertension:** Summary Table: <http://www.rxfiles.ca/rxfiles/uploads/documents/HTNandmarkHypertensionTrials.pdf>  
ACCOMPLISH: <http://www.rxfiles.ca/rxfiles/uploads/documents/HTN-QandA-Accomplish.pdf>  
ALLHAT: <http://www.rxfiles.ca/rxfiles/uploads/documents/HTN-Update-2003-Final.pdf>  
ANBP2: <http://www.rxfiles.ca/rxfiles/uploads/documents/HTN-QandA-ANBP2.pdf>  
ASCOT-BPLA: <http://www.rxfiles.ca/rxfiles/uploads/documents/HTN-QandA-ASCOT.pdf>  
SPRINT: <http://www.rxfiles.ca/rxfiles/uploads/documents/SPRINT-BP-Trial-Overview.pdf> <sup>2015</sup>  
Trial Summary table - abridged: <http://www.rxfiles.ca/rxfiles/uploads/documents/members/cht-HTN-trial-summary.pdf>  
**HF:** CHARM: <http://www.rxfiles.ca/rxfiles/uploads/documents/CHARM-Comments.pdf>  
PARADIGM-HF: <http://www.rxfiles.ca/rxfiles/uploads/documents/PARADIGM-HF-Trial-Sacubitril.pdf> <sup>2015</sup>  
**Hirsutism:** <http://www.rxfiles.ca/rxfiles/uploads/documents/members/Hirsutism%20Trial%20Summary.pdf>  
**HRT:** WHI: <http://www.rxfiles.ca/rxfiles/uploads/documents/HRT-Post-WHI-2002-Header.pdf>  
WHI & Age: <http://www.rxfiles.ca/rxfiles/uploads/documents/HRT-Age-and-the-WHI.pdf> ;  
WHI & Extras/Perspectives on NNTs, NNHs: <http://www.rxfiles.ca/rxfiles/uploads/documents/HRT-WHI-Extras-Perspectives.pdf>

**Lipid:** Summary Table: [http://www.rxfiles.ca/rxfiles/uploads/documents/members/CHT-lipid\\_agents-major\\_trials.pdf](http://www.rxfiles.ca/rxfiles/uploads/documents/members/CHT-lipid_agents-major_trials.pdf)  
& Q&A 2004: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-QandA-Update-Oct04.pdf>  
AIM-HIGH: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-AIM-HIGH-nicotinic-acid-Niaspan-trial.pdf>  
ASCOT-LLA: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-QandA-ASCOT.pdf>  
CARDS: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-QandA-CARDS.pdf>  
ENHANCE: [http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-ENHANCE\\_trial\\_overview.pdf](http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-ENHANCE_trial_overview.pdf)  
FIELD Substudy: <http://www.rxfiles.ca/rxfiles/uploads/documents/FIELD-Sub-Analysis-Women-Trial-Summary.pdf> <sup>2015</sup>  
IDEAL: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-QandA-IDEAL.pdf>  
IMPROVE-IT: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-IMPROVE-IT-Trial-Summary-QandA.pdf> <sup>2014</sup>  
JUPITER: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-Jupiter-trial-overview.pdf>  
PROVE-IT: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-QandA-Prove-It.pdf>  
SHARP: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-Sharp-CKD-trial.pdf>  
SPARCL: <http://www.rxfiles.ca/rxfiles/uploads/documents/Lipid-QandA-SPARCL.pdf>  
**Thrombotic (antithrombotics: ASA, clopidogrel, anticoagulants: warfarin) :**  
ACTIVE-A & ACTIVE-W trials <http://www.rxfiles.ca/rxfiles/uploads/documents/ACTIVE-A-Trial-Summary.pdf>  
Antithrombotics Summary Chart: <http://www.rxfiles.ca/rxfiles/uploads/documents/members/cht-AntiThrombotics.pdf>  
ARISTOTLE: Apixaban vs warfarin in A Fib: <http://www.rxfiles.ca/rxfiles/uploads/documents/ARISTOTLE-AF-Apixaban.pdf>  
CHARISMA: <http://www.rxfiles.ca/rxfiles/uploads/documents/Charisma-QandA.pdf>  
Clopidogrel-PPI drug interaction: <http://www.rxfiles.ca/rxfiles/uploads/documents/Clopidogrel-PPI-interaction-QandA.pdf>  
DAPT: 12 vs 30months <http://www.rxfiles.ca/rxfiles/uploads/documents/DAPT-Trial-12vs30months.pdf>  
PCI-Clarity: <http://www.rxfiles.ca/rxfiles/uploads/documents/PCI-CLARITY%20Trial%20Summary.pdf> <sup>2016</sup>  
PCI-CURE: <http://www.rxfiles.ca/rxfiles/uploads/documents/PCI-CURE%20Trial%20Summary.pdf> <sup>2016</sup>  
PEGASUS-TIMI 54: Ticagrelor vs P1, prior-MI: <http://www.rxfiles.ca/rxfiles/uploads/documents/PEGASUS%20Trial%20Summary.pdf> <sup>2016</sup>  
PLATO: Ticagrelor vs clopidogrel ACS: <http://www.rxfiles.ca/rxfiles/uploads/documents/PLATO%20Trial%20Summary.pdf> <sup>2016</sup>  
RE-LY: Dabigatran vs warfarin in Atrial Fibrillation <http://www.rxfiles.ca/rxfiles/uploads/documents/RE-LY-Trial-Dabigatran.pdf> <sup>2016</sup>  
ROCKET-AF: Rivaroxaban vs warfarin in A Fib: <http://www.rxfiles.ca/rxfiles/uploads/documents/ROCKET-AF-Rivaroxaban.pdf>  
TRITON-TIMI 38: Prasugrel vs clopidogrel, ACS: <http://www.rxfiles.ca/rxfiles/uploads/documents/TRITON-TIMI%2038%20Trial%20Summary.pdf> <sup>2016</sup>  
**MISC.:**  
Catie-AD: Atypical Antipsychotics in Patients with Alzheimer's <http://www.rxfiles.ca/rxfiles/uploads/documents/Psych-CATIE-AD-trial-summary.pdf>  
FLAME: Indacaterol+Glycopyrronium vs Salmeterol+Fluticasone for COPD: <http://www.rxfiles.ca/rxfiles/uploads/documents/FLAME-Trial-Summary.pdf> <sup>2016</sup>  
Meloxicam: SELECT, MELISSA; celecoxib CLASS, rofecoxib VIGOR. : <http://www.rxfiles.ca/rxfiles/uploads/documents/QandA-Meloxicam-2.pdf>  
OAB: Darifenacin-Oxybutynin Memory Trial : <http://www.rxfiles.ca/rxfiles/uploads/documents/UI-Darifenacin-Kay-Trial-QandA.pdf>  
SENIOR: <http://www.rxfiles.ca/rxfiles/uploads/documents/Senior-Trial-Oxybutynin-Solifenacin-Elderly-Cognitive-Impairment.pdf>  
WARFASA: <http://www.rxfiles.ca/rxfiles/uploads/documents/Aspirin-warfarin-trial-summary-WarfASA.pdf>

## Online Extras

### Guidelines, Evidence and Patients

- Guidelines provide guidance based on evidence, clinical experience and someone's values and preferences.
  - Considerations: Evidence
    - How current? Has level/strength of evidence (evidence quality) been assessed for recommendations? (e.g. May have more confidence if high quality, multiple, trials.)
    - What outcomes are evaluated? Are they patient orientated or surrogate?
    - Has evidence been allowed to inform pre-existing assumptions, biases and beliefs?
    - Is the evidence applicable to your patient(s)?
  - Clinical experience / consensus
    - Is expert opinion, and extent of agreement acknowledged
  - Conflicts of Interest (<https://www.cmaj.ca/content/193/2/E49>)
    - Are conflicts of interest disclosed
    - Conflicts may be financial or non-financial
    - Methods should be transparent and rigorous to inform objectively on best available evidence
  - Values
    - In what way are values and preferences included?
    - Who's values: Patient? Society? Payer? Professional?

- Overall assessment:
  - Look for transparency, evidence ratings, peer review, conflicts of interest
  - Do the guidelines allow for, and enable, shared decision making with patients?
- If the guidelines don't apply, don't apply them! Almost all guidelines contain a chapter/disclaimer noting that any recommendations must be assessed and individualized for the patient in front of you. Recommendations are often intended to apply to a majority of patients, but may not be suitable for the patient in front of you. If so, document reason for your decision.

EBM	1
Evidence Based Medicine	1
NNH	1
NNT	1
Precision	1
Validity	1

1 Sackett D, Straus S, Richardson WS, Rosenberg, Haynes R. Evidence-Based Medicine: how to practice and teach EBM. Churchill Livingstone. 2000.

2 Davies M, Davies P, Gray A, Mant J, Seers K, Snowball. Evidence-based Practice: a primer for health care professionals. Elsevier, 2<sup>nd</sup> edition 2005.

3 Allen J. Pharmacist's Letter / Prescriber's Letter. Applying Study Results to Patient Care. June 2005, 21:1-14.

4 Jadad A, Enkin M. Randomized Controlled Trials: Questions, Answers and Musings 2<sup>nd</sup> edition. Blackwell Publishing 2007; BMJ Books

5 Edited by: Geyman J, Deyo R, Ramsey S. Evidence-Based Clinical Practice: Concepts and Approaches. Butterworth Heinemann 2000.

6 Centre for Evidence Based Medicine (CEBM) Oxford: EBM Tools accessed online 31Jul08 at: <http://www.cebm.net/index.aspx?o=1157>

7 Centre for Evidence Based Medicine, University Health Network, Toronto. Critical Appraisal tools; accessed online 31Jul08: <http://www.cebm.utoronto.ca/teach/materials/caworksheets.htm>

8 Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. JAMA. 1995 Feb 1;273(5):408-12. [Compared with trials in which authors reported adequately concealed treatment allocation, trials in which concealment was either inadequate or unclear (did not report or incompletely reported a concealment approach) yielded larger estimates of treatment effects (P < .001). Odds ratios were exaggerated by 41% for inadequately concealed trials and by 30% for unclearly concealed trials (adjusted for other aspects of quality). Trials in which participants had been excluded after randomization did not yield larger estimates of effects, but that lack of association may be due to incomplete reporting. Trials that were not double-blind also yielded larger estimates of effects (P = .01), with odds ratios being exaggerated by 17%.]

9 Moher D, Pham B, Jones A, Cook DJ, Jadad AR, Moher M, Tugwell P, Klassen TP. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? Lancet. 1998 Aug 22;352(9128):609-13. [FINDINGS: The quality of trials was low. Masked assessments provided significantly higher scores than unmasked assessments (mean 2.74 [SD 1.10] vs 2.55 [1.20]). Low-quality trials (score < or = 2), compared with high-quality trials (score > 2), were associated with an increased estimate of benefit of 34% (ratio of odds ratios [ROR] 0.66 [95% CI 0.52-0.83]). Trials that used inadequate allocation concealment, compared with those that used adequate methods, were also associated with an increased estimate of benefit (37%; ROR=0.63 [0.45-0.88]). The average treatment benefit was 39% (odds ratio [OR] 0.61 [0.57-0.65]) for all trials, 52% (OR 0.48 [0.43-0.54]) for low-quality trials, and 29% (OR 0.71 [0.65-0.77]) for high-quality trials. Use of all the trial scores as quality weights reduced the effects to 35% (OR 0.65 [0.59-0.71]) and resulted in the least statistical heterogeneity. INTERPRETATION: Studies of low methodological quality in which the estimate of quality is incorporated into the meta-analyses can alter the interpretation of the benefit of intervention, whether a scale or component approach is used in the assessment of trial quality.]

10 Fletcher J. Subgroup analyses: how to avoid being misled. BMJ. 2007 Jul 14;335(7610):96-7.

11 Centre for Evidence Based Medicine (CEBM) Oxford: EBM Tools accessed online 31Jul08 at: <http://www.cebm.net/index.aspx?o=1039>

12 Oleckno WA. Essential Epidemiology Principles and Applications. Long Grove IL: Waveland Press Inc. 2002. (page 108)

13 <http://www.cochrane.org/>; <http://www.cfp.ca/cgi/content/full/55/11/1155?etoc>

14 <http://www.campbellcollaboration.org/>; <http://www.campbellcollaboration.org/library.php>

15 Centre for Evidence Based Medicine (CEBM) Oxford: EBM Tools accessed online 31Jul08 at: <http://www.cebm.net/index.aspx?o=1025>

16 Barratt A, Wyer PC, Hatala R, McGinn T, Dans AL, Keitz S, Moyer V, For GG. Evidence-Based Medicine Teaching Tips Working Group. Tips for learners of evidence-based medicine: 1. Relative risk reduction, absolute risk reduction and number needed to treat. CMAJ. 2004 Aug 17;171(4):353-8.

17 Bandler: Number Needed to Treat. Accessed online 31Jul08 at: <http://www.medicines.ox.ac.uk/bandler/band59/NNT1.html>

18 Finlay A, McAlister. The "number needed to treat" turns 20 — and continues to be used and misused. CMAJ 2008; 179: 549-553. <http://www.cmaj.ca/cgi/content/full/179/6/549>

19 James P McCormack. NNT misses the mark on baseline risks. Electronic Letter, CMAJ online. (17 September 2008) <http://www.cmaj.ca/cgi/eletters/179/6/549#20357>

20 Centre for Evidence Based Medicine (CEBM) Oxford: NNT calculator tool. Accessed online 31Jul08 at <http://www.cebm.net/index.aspx?o=1044>.

21 Montori VM, Kleinbart J, Newman TB, Keitz S, Wyer PC, Moyer V, Guyatt G. Evidence-Based Medicine Teaching Tips Working Group. Tips for learners of evidence-based medicine: 2. Measures of precision (confidence intervals). CMAJ. 2004 Sep 14;171(6):611-5.

22 Hatala R, Keitz S, Wyer P, Guyatt G. Evidence-Based Medicine Teaching Tips Working Group. Tips for learners of evidence-based medicine: 4. Assessing heterogeneity of primary studies in systematic reviews and whether to combine their results. CMAJ. 2005 Mar 1;172(5):661-5.

23 Kadoszkiewicz H, Zimmermann T, Beck-Bornholdt HP, van den Busche H. Cholinesterase inhibitors for patients with Alzheimer's disease: systematic review of randomised clinical trials. BMJ. 2005 Aug 6;331(7512):321-7. <http://www.bmj.com/cgi/rapidprint/331/7512/321> [A systematic review by Kadoszkiewicz and colleagues (p 321) included 22 double blind randomised controlled trials with the follow-up ranging from six weeks to three years, but the trials scored poorly on a predefined checklist of criteria of methodological quality. Further, the outcomes measuring cognition did show beneficial effects of cholinesterase inhibitors, but these effects were minimal (ranging from 1.5 points to 3.9 points on a 70 point Alzheimer's disease assessment scale).]

24 Regier L. RxFiles Trial Summary: Darifenacin (ENABLEX) vs Oxybutynin ER <sup>extended release</sup> (DITROPAN XL) vs Placebo: Effects On Memory / Cognitive Impairment. Accessed at: <http://www.rxfiles.ca/rxfiles/uploads/documents/UI-Darifenacin-Kay-Trial-QandA.pdf>

25 Montori VM, Isley WL, Guyatt GH. Waking up from the DREAM of preventing diabetes with drugs. BMJ. 2007 Apr 28;334(7599):882-4.

26 Montori VM. Treat the Target or Treat the Patient. Aust Prescr 2011;34:94-5. Accessed online Aug 2, 2011 <http://www.austliianprescriber.com/magazine/34/4/94/5>.

27 Halvorsen PA, Selmer R, Kristiansen IS. Different ways to describe the benefits of risk-reducing treatments: a randomized trial. Ann Intern Med. 2007 Jun 19;146(12):848-56. Summary for patients in: Ann Intern Med. 2007 Jun 19;146(12):i50.

28 Anonymous. Intensive blood glucose control with sulphonylureas or insulin compared with conventional treatment and the risk of complications in patients with type 2 diabetes (UKPDS 33). Lancet 1998;352:837-53.

29 Regier L. HRT in Light of the WHI — Data in Perspective. Sept 2002. Available online at: <http://www.rxfiles.ca/rxfiles/uploads/documents/HRT-WHI-Extras-Perspectives.pdf>

30 Ross JS, Mulvey GK, Hines EM, et al. Trial publication after registration in ClinicalTrials.gov: A cross-sectional analysis. PLoS Med 2009; DOI:10.1371/journal.pmed.1000144.

31 Mathieu S, Boutron I, Moher D, et al. Comparison of registered and published primary outcomes in randomized controlled trials. JAMA 2009; 302: 977-984.

32 McNutt Robert A.; Livingston Edward H. Evidence-Based Medicine Requires Appropriate Clinical Context. JAMA. 2010; 303(5):454-455.

33 Kent DM, Steyerberg E, van Klaveren D. Personalized evidence based medicine: predictive approaches to heterogeneous treatment effects. BMJ. 2018 Dec 10;363:k4245.

34 Laffin LJ, Besser SA, Alenghat FJ. A data-zone scoring system to assess the generalizability of clinical trial results to individual patients. Eur J Prev Cardiol.2018;2047487318815967.

35 Senn SJ, Lewis RJ. Treatment Effects in Multicenter Randomized Clinical Trials. JAMA. 2019 Mar 1.

36 CADTH: <https://www.cadth.ca/>

## RxFiles: Evidence Based Medicine (EBM) Overview - References

**Original Reviewer Acknowledgments:** G Michael Allen MD, CCFP, Associate Professor, Director of EBM, Dept of Fam Med, U of A. Michael Allen MD, Associate Professor, Director Evidence-based Programs, Dalhousie University CME, Pam Maclean-Veysey, Drug Eval Unit, Halifax. Derek Jorgenson, PharmD, U of S. Darcy Lamb, C of Pharmacy, U of S. David Blackburn, C of Pharmacy, U of S. G. Turcotte MD, FM/EBM teaching, Gatineau, Quebec. Loren D Regier www.RxFiles.ca

37 Falk J, Thomas B, Kirkwood J, Korowynk CS, Lindblad AJ, Ton J, et al. PEER systematic review of randomized controlled trials: Management of chronic neuropathic pain in primary care. Can Fam Physician. 2021 May;67(5):e130-e140. Accessed online 21May21 at: <https://www.cfp.ca/content/67/5/e130.long>

38 Frieden TR. Evidence for Health Decision Making - Beyond Randomized, Controlled Trials. N Engl J Med. 2017 Aug 3;377(5):465-475.

39 Djulbegovic B. Ethics of uncertainty. Patient Educ Couns. 2021 Nov;104(11):2628-2634. doi: 10.1016/j.pec.2021.07.025. Epub 2021 Jul 15. PMID: 34312034. [Accessed online](https://doi.org/10.1016/j.pec.2021.07.025), 15Nov2021.

40 Schandelmaier S, Briel M, Varadhan R, Sauerbrei W, Devasenapathy N, Hayward RA, Gagnier J, Borenstein M, VanderHeijden GJMG, Dahabreh I, Schmid CH, Ioannidis JPA, Walsh M, Thabane L, Guyatt GH. Development of a new Instrument to assess the Credibility of Effect Modification Analyses (ICEMAN) in randomized controlled trials and meta-analyses. CMAJ. 2020 192(32):E901-E906

## Additional References:

Alahdab F, Murad MH. Evidence maps: a tool to guide research agenda setting. BMJ Evid Based Med. 2019 Dec;24(6):209-211.

Aubin D, Hebert M, Eurich D. The importance of measuring the impact of patient-oriented research. CMAJ. 2019 Aug 6;191(31):E860-E864.

Bibens M, Vassar M, Wayant C. Use of a meta-research team to facilitate evidence-based medicine to the next generation. BMJ Evid Based Med. 2019 Dec;24(6):205-206.

Brassey J, Price C, Edwards J, et al. Developing a fully automated evidence synthesis tool for identifying, assessing and collating the evidence. BMJ Evid Based Med. 2019 Aug 29.

Chiu B, Liu M, Leas EC, et al. Effect size reporting among prominent health journals: a case study of odds ratios. BMJ Evidence-Based Medicine Epub ahead of print: Aug 2021. doi:10.1136/bmjebm-2020-111569. Accessed 04 Aug 2021 online at <https://ebm.bmj.com/content/ebm/2020/12/10/bmjebm-2020-111569.full.pdf>

Da-Ré R. Analysis of retracted articles on medicines administered to humans. Br J Clin Pharmacol. 2019 Jun 24.

Frank O, Tam CM, Rhee J. Is it time to stop using statistical significance? Aust Prescr. 2021 Feb;44(1):16-18. doi: 10.18773/austprescr.2020.074. Epub 2021 Feb 1. PMID: 33664545; PMCID: PMC7900272. Accessed online 04 Aug 2021 at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7900272/>

Ghanmad M, Yang B, Leeftang M, Aldroft A, Bossuyt PM, Schroter S, Boutron I. A randomized trial of an editorial intervention to reduce spin in the abstract's conclusion of manuscripts showed no significant effect. J Clin Epidemiol. 2021 Feb;130:69-77. doi: 10.1016/j.jclinepi.2020.10.014. Epub 2020 Oct 21. PMID: 33096222. [Related: An unsuccessful initiative to reduce 'spin' in biomedical literature. Drug Ther Bull. 2021 Dec;59(12):182. doi: 10.1136/dtb.2021.000059. Epub 2021 Nov 12. PMID: 34772742.]

Goupil B, Balusson F, Naudet F, et al. Association between gifts from pharmaceutical companies to French general practitioners and their drug prescribing patterns in 2016: retrospective study using the French Transparency in Healthcare and National Health Data System databases. BMJ. 2019 Nov 5;367:i6015.

Herrera-Perez D, Haslam A, Crain T, et al. A comprehensive review of randomized clinical trials in three medical journals reveals 396 medical reversals. Elife. 2019 Jun 11;8.

Ioannidis JPA. Options for publishing research without any P-values. Eur Heart J. 2019 Aug 14;40(31):2555-2556.

John LK, Loewenstein G, Marder A, Callahan ML. Effect of revealing authors' conflicts of interests in peer review: randomized controlled trial. BMJ. 2019 Nov 6;367:i5896.

Khan MS, Lateef N, Siddiqi TJ, et al. Level and Prevalence of Spin in Published Cardiovascular Randomized Clinical Trial Reports With Statistically Nonsignificant Primary Outcomes: A Systematic Review. JAMA Netw Open. 2019 May 3;2(5):e192622.

Messerli FH, Bangalore S, Messerli AW. Meta-Analysis in the Mirror of its Quotations: Science, Scepticism, Scorn, and Sarcasm. Eur Heart J. 2019 Oct 21;40(40):3290-3291.

Mostofsky E, Dunn JA, Hernández-Díaz S, Mittleman MA. Patient and Physician Preferences for Reporting Research Findings. Fam Med. 2019 Jun;51(6):502-508.

Murray SB, Heathers JA, Schauer RM, et al. Postpublication Metrics of Randomized Clinical Trials With and Without Null Findings. JAMA. 2019 May 14;321(18):1825-1826.

Niforatos JD, Weaver M, Johansen ME. Assessment of Publication Trends of Systematic Reviews and Randomized Clinical Trials, 1995 to 2017. JAMA Intern Med. 2019 Jul 29.

Nissen SE, Reed GW. Can we trust observational data for clinical decision-making? Eur Heart J. 2018 Dec 27.

Norton EC, Dowd BE, Maciejewski ML. Marginal Effects-Quantifying the Effect of Changes in Risk Factors in Logistic Regression Models. JAMA. 2019 Mar 8.

Park JH, Detry MA, Murthy S, Guyatt G, Mills EJ. How to Use and Interpret the Results of a Platform Trial: Users' Guide to the Medical Literature. JAMA. 2022 Jan 4;327(1):67-74.

Reynolds-Vaughn V, Riddle J, Brown J, et al. Evaluation of Spin in the Abstracts of Emergency Medicine Randomized Controlled Trials. Ann Emerg Med. 2019 May 14.

Sanders GD, Maciejewski ML, Basu A. Overview of Cost-effectiveness Analysis. JAMA. 2019 Mar 11.

Sterne JAC, Savovic J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. BMJ. 2019 Aug 28;366:i4898.

Trinquart L, Erlinger AL, Petersen JM, et al. Applying the E-value to Assess the Robustness of Epidemiologic Fields of Inquiry to Unmeasured Confounding. Am J Epidemiol. 2019 Mar 15.

Wilson FP. Free yourself from study fragility and P-values. Medscape. Sept 25, 2019. Accessed online 21 Oct 2019 at <https://www.medscape.com/viewarticle/918733>.

Yao L, Ahmed MM, Guyatt GH, et al. Discordant and inappropriate discordant recommendations in consensus and evidence based guidelines: empirical analysis. BMJ. 2021 Nov 25;375:e066045.